

THERAPEUTIC POTENTIAL AND PHYTOCHEMISTRY OF NATURAL HERBALS OF FAMILY LAMIACEAE: *OCIMUM SANCTUM* L. AND *OCIMUM BASILICUM*

Lata Rana

Department of Chemistry, D. S. B. Campus, Kumaun University, Nainital 263001, India

Geeta Tewari*

Department of Chemistry, D. S. B. Campus, Kumaun University, Nainital 263001, India

Deepshrekha Punetha

Delma Laboratory for Soil & Construction Material Inspection L. L. C. Abu Dhabi (United Arab Emirates)

Mamta Bisht

*Department of Chemistry, D. S. B. Campus, Kumaun University, Nainital 263001, India
Department of Chemistry, Uttarakhand University, Dehradun, India*

Geetanjali Upadhyay

Department of Botany, D. S. B. Campus, Kumaun University, Nainital 263001, India

Chitra Pande

Department of Chemistry, D. S. B. Campus, Kumaun University, Nainital 263001, India

Lalit M. Tewari

Department of Botany, D. S. B. Campus, Kumaun University, Nainital 263001, India

Keywords: Biodiversity, Lamiaceae, *Ocimum sanctum*, *Ocimum basilicum*, anticancer, nanomedicine, nutraceuticals, aromaceuticals, antiviral, anti-inflammatory, anti-diabetic, insecticidal

Contents

1. General Introduction
2. History behind Ocimum
 - 2.1. In Ayurveda
 - 2.2. In Greek-Unani system
 - 2.3. In Homeopathy
 - 2.4. In Tibetan Medicine Systems
3. The Lamiaceae Family
 - 3.1. List of Some Ocimum Species Found in India
 - 3.2. Synonyms and Common Name
 - 3.3. Importance of Ocimum Genus
 - 3.4. Taxonomic Classification of Basil (*Ocimum basilicum*) and Tulsi (*Ocimum sanctum*)
 - 3.5. Description of Basil (*Ocimum basilicum*) and Tulsi (*Ocimum sanctum*)
 - 3.6. Distribution of Basil (*Ocimum basilicum*) and Tulsi (*Ocimum sanctum*)
4. Climatic Conditions for Plant Growth

5. Phytochemicals Present in Basil and Tulsi

5.1. Extract of Tulsi (*Ocimum sanctum*) and Basil (*Ocimum basilicum*)

5.2. Essential Oils of Tulsi (*Ocimum sanctum*) and Basil (*Ocimum basilicum*)

6. Biological Activities (as pharmaceuticals) of Plant Extracts of Tulsi (*Ocimum sanctum*) and Basil (*Ocimum basilicum*)

6.1. Antioxidant Activity

6.2. Antimicrobial Activity

6.3. Anti-inflammatory Activity

6.4. Anticancer Activity

6.5. Anti-diabetic activity

6.6. Antiviral Activity

6.7. Insecticidal Activity

6.8. Miscellaneous Activities

7. Biological Properties (as Pharmaceuticals) of Essential Oil of Tulsi (*O. sanctum*) and Basil (*O. basilicum*)

7.1. Antioxidant Activity

7.2. Antimicrobial Activity

7.3. Anti-inflammatory Activity

7.4. Anti-cancer Activity

7.5. Anti-diabetic Activity

7.6. Antiviral Activity

7.7. Insecticidal Activity

7.8. Miscellaneous Activities

8. Other Applications (Herbal Based Products for Health Care and Personal Care)

8.1. Aroma Therapy (Aromaceuticals)

8.2. Cosmetics (Cosmeceuticals)

8.3. Disinfectants

8.4. Food Packaging and Storage

8.5. Nutrition Supplement (Neutraceuticals)

8.6. Perfumery

8.7. Repellents

9. Medicinal Benefits

9.1. Common Cough, Colds and Fever

9.2. Headache and Stress

9.3. Inflammation

9.4. Open Wounds

9.5. Diarrhea

9.6. Skin Disorders

9.7. Nanomedicine

9.8. Others

10. Future Perspectives

11. Conclusion

Glossary

Bibliography

Biographical Sketches

Summary

Rich biodiversity of natural products offers a vast area for phytochemical and pharmacological investigations. Therefore, at present over 80% of known (approximately 30,000) natural products have originated from plants worldwide. Amongst some of the astonishing herbs, family Labiate (syn. Lamiaceae) includes more than 252 genera and 7000 species. The genus *Ocimum* is one of the extensively used member of Lamiaceae family and comprising 150 species. This study elaborated the vast global applicability of *Ocimum basilicum* and *Ocimum sanctum* extracts as well as their essential oils in recent markets as important components in organic cosmetics, perfumery, food packaging, storages, nutraceuticals, ethnomedicinal, therapeutic and pharmacology. Moreover, *O. basilicum* and *O. sanctum* could be an important valuable source of substances, which possesses remarkable antioxidant, antibacterial, antifungal, anti-hepatotoxicity, anticancer, antiviral, antidiabetic, antimicrobial, anti-inflammatory, analgesic and various other activities. Thus, a systematic review was performed to investigate the phytochemical and pharmacological potential of *O. sanctum* and *O. basilicum* to search the possibilities of these metabolites in the health sector for discovery and development of new drugs. Additionally, essential oils of *O. sanctum* and *O. basilicum* were developed for the determination of various micro and nano herbal drug substances via nanotechnology. They also have important roles in emerging development of herbal cosmeceuticals which reduces the risk of side effects of some synthetic chemicals.

In this context, the proposed chapter is the outcome of various collaborative works from all over the world, which includes a brief introduction along with historical content. Thereafter, phytochemistry and biological potential of extracts as well as essential oils of both species will be discussed. Ending section of this chapter highlights the advantages, conclusions and future prospectives.

1. General Introduction

Despite the various advancements in diagnosis and surgery, today's world is fretting with various chronic diseases. Recently, an infectious disease COVID-19 caused by corona virus, is an example of that how the world is suffering with newly emerging diseases. So, the priority in the present times is to build an adequate immune system to fight against diseases which is a big challenge for scientists and researchers across the world. Strong immune system plays a powerful role in the prevention of infections or viruses. During the past few years, the interest in the natural products-based medication has considerably increased for strengthening the immunity because of their mild to no side effects. Natural products are rich and wide sources of potentially active secondary metabolites such as terpenoids, alkaloids, and flavonoids. These secondary metabolites contain considerable range of antioxidant, antimicrobial, antiviral, anti-inflammatory, anti-diarrheal, anti-diabetic, cardio protective, analgesic, antipyretic, anti-tumor, anti-stress, anti-allergic, antihypertensive, anticancer and many more activities. Moreover, approximately 13000 plant species have been found which are being used as drug throughout the globe.

The use of natural products for sustainable development has been explored globally in few last decades. Among the species known for therapeutic potential, the aromatic species and their essential oils are more commercially valuable. Essential oils are also a natural source of antioxidants and widely acceptable by the consumers. Nowadays, principally active components of essential oil extracted from aromatic plant species are being utilized in various clinical applications, food and flavorings industries, cosmetics and perfume markets. Over the last decades, the interest in essential oil and its trends in beauty products have grown significantly. Generally, essential oils are the mixture of volatile compounds, which show low stability towards heat, oxygen and moisture. Therefore, various new techniques have been developed to prevent them from degradations. Drying is one of the promising techniques which removes contamination and increases self-life of the herbal products. Recently, natural products like herbal tea, soup, syrup and decoction are vastly consumed and drying technique could be another way of increasing consumption of natural based products for long time. Previous studies reported that drying of herbal species can improve their bio activities and potential for various sicknesses. Nano- and micro- encapsulation of the essential oil is the most convenient method to protect it from microorganism, fungus and enhances direct or indirect uses of essential oil to cure various human ailments.

Some of the most important and popular aromatic herbs are lavender, rosemary, peppermint, sage, fennel, artemisia chameleon, oregano, coriander, red spider lily, thyme, geranium also including the king of herbs '*Ocimum*'. *Ocimum* has been famous as royal herb in French and sacred herb in India. In Hinduism, *Ocimum* (Tulsi) has great spiritual importance and used in worship (Figure 1).




Figure 1. Phytochemical and pharmacological potential of *O. sanctum* L. and *O. basilicum* L.

2. History behind *Ocimum*

Ocimum commonly known as Tulsi, which is mentioned in the ancient language of Sanskrit as Tulasi, Ajaka, Krishnamul and its translated meaning is “the matchless one”. In Arabic, *Ocimum* is called by the name Raihaan and Habaq. Moreover, in French language, it is popular as Basilic Saint, In Hindu scripture, according to *Padma Purana*, importance of *Ocimum* (Tulsi) in worship was described and mentioned in a chapter of Tulsi Vivah. Tulsi is considered as a goddess and consort of lord Vishnu. Therefore, *Ocimum* (Tulsi) is also called “*Vishnupriya*”. In India, some morphotypes of *Ocimum* species, also known as “Rama or Shri or Lakshmi Tulsi” and “Shyama or Krishna Tulsi” according to their leave’s color; the light green and dark purplish respectively. *Ocimum basilicum* is also known by various names such as Basil/Common Basil or Sweet Basil in English, whereas, it is called as *Babui Tulsi* in Hindi and Bengali. The plant is known as *Badrooj*, *Hebak* or *Rihan* in Arabic; as *Nasabo* or *Sabje* in Gujrati and as *Jangli Tulsi* in Urdu. *Tohrakhurasani* and *Okimon* are the other ascribed names.

2.1. In Ayurveda

Importance of Tulsi in Indian traditional therapeutics system with ancient theories has been indicated in the *Charaka Samhita* (an ancient Sanskrit text on Ayurveda). In Ayurveda, Tulsi has been considered as a medicinal herb and a source of strong aroma containing components for several decades. Its diverse healing potential along with stress relief properties has been earlier mentioned. Tulsi extracts are used to cure common cold, fever, cough, stomach infections and many more.

2.2. In Greek-Unani system

Unani medicine (Yunani) is known for its wide range of adaptability in Muslim culture mostly in South Asia and more recently, in some other parts of Asian countries. Unani medicine is commonly called as Perso-Arabic traditional medicine system and pseudoscientific system. In Unani aromatic herbal system, more than 1000 plants are still in use including Faranjmishk (*Ocimum gratissimum* Linn.) with many folk applications in nostrils, obstruction of brain, common fever etc. Basil was used to produce royal perfumes in Greece, hence it was named as “royal plant”.

2.3. In Homeopathy

Among the herbs known for medicinal value, *Ocimum* has wide history of uses in traditional Homoeopathy system. The efficacy of Homoeopathic medicines depends upon the active compounds present in the herbal medicinal plants. Various species of *Ocimum* are major sources of these bioactive components. Therefore, in terms of easy accessibility, high effectiveness, mild to no side effects and economical aspects, the low cost homoeopathic *Ocimum* mother tincture (a mixture or extract of herbal plant) is a good alternative for the primary health care and 70-80% of the world’s population uses homoeopathic medicine in day-to-day practices.

2.4. In Tibetan Medicine Systems

For a long time, more than 2000 years, Traditional Tibetan medicine system (formerly called Sowa-Rigpa medicine) had a wide and fanciful history including various unique techniques such as pulse readings, physical therapies, urinalysis and preferred natural herbs to treat illness. In addition, latest statistics suggested that the Tibetan medical system is based upon 2,644 natural plants including *Ocimum* and a total of 3,105 natural medicines are being practiced in daily life. Huge quantities of Tibetan medicines are used in India, Tibet, Ladakh, Nepal, Bhutan, China, Mongolia and Siberia also more recently in many parts of Europe, North America and some other parts of the world.

3. The Lamiaceae Family

Family Lamiaceae (Labiatae), is one of the most important essential oil-bearing families of more than 252 genera and 7000 species. Since ancient, Lamiaceae family is also called as mint family of flowering plants and majorities of them are aromatic, herbage with clinical properties. Best known species of the largest family includes *Lavender*, *Ocimum*, *Rosemary*, *Origanum*, *Hyssop*, *Thymus*, *Sage*, *Mentha*, *Nepeta*, *Perilla* etc. Aromatic species in the family are rich sources of naturally occurring antioxidants and used in various pharmaceuticals, cosmetics, perfumery, food flavoring agents and many other industrial applications. Even though, most of the species are generally cultivated in home gardens for their fresh scent leaves and interesting flowers. Intense fragrance of these aromatic species is also used in aromatherapy. Many of the species of family Lamiaceae which contain essential oil have attained largest economical value in recent years. Their essential oil play major role in cosmetics and perfume market and demand of essential oil is increasing day by day. The well-known *Ocimum* of this family singly composed of 150 species, approximately most of which are aromatic and still in use.

3.1. List of Some *Ocimum* Species Found in India

Ocimum sanctum L.
Ocimum basilicum L.
Ocimum gratissimum L.
Ocimum americanum L.
Ocimum kilimandschricum Gürke
Ocimum filamentosum Forssk.
Ocimum minimum L.

Figure 2 represents the photographs of (a) *O. sanctum* (b) *O. basilicum* (c) *O. gratissimum* (d) *O. americanum*

3.2. Synonyms and Common Name

Ocimum sanctum L.; syn. *O. tenuiflorum* (Holy basil/Tulsi)
Ocimum basilicum L.; syns. *O. album* L., *O. anisatum* Benth. (Basil)
Ocimum gratissimum L. *Ocimum viride* Willd. (Clove basil/African basil)
Ocimum americanum L. (*O. canum* Sims) (Hoary basil)

Ocimum kilimandschricum Gürke (African blue basil)
Ocimum filamentosum Forssk.; syn. *O. adscendens* Willd.
Ocimum minimum L. (Bush basil)




Figure 2. The photographs of (a) *O. sanctum* (b) *O. basilicum* (c) *O. gratissimum* (d) *O. americanum* in natural environment

3.3. Importance of *Ocimum* Genus

The genus *Ocimum* is one of the well-known genera in the Lamiaceae (mints) family. The genus is a big provider of more than 150 species globally. Majority of the *Ocimum* species are aromatic and medicinally valuable. Presence of potentially active compositions has been mentioned in several species of *Ocimum*. It is an important species for research purpose because of its versatile applications, cost-effectiveness, and ease of availability. In this novel pandemic period, when most of the countries in the world are suffering from corona virus, Tulsi is used as an active integrant of homemade decoction (*kadha*) and soup against corona virus. In Ayurvedic studies, Tulsi has already been mentioned as an immune booster. It has been reported to exhibit analgesic, anti-diabetic, anti-inflammatory, antimicrobial, anti-tumour, anti-stress, anti-viral, cardioprotective, immunomodulatory, neuroprotective and rejuvenating potential. On the other hand, dried plant parts of this species are consumed in the form of herbal tea in day-to-day basis. Spices of *Ocimum* are commonly used in various dishes, soup and syrups. Basil (*Ocimum basilicum*) and Tulsi (*Ocimum sanctum*) are two most important aromatic herbs of genus *Ocimum*.

3.4. Taxonomic Classification of Basil (*Ocimum basilicum*) and Tulsi (*Ocimum sanctum*)

Figure 3 illustrates the taxonomical position of two *Ocimum* species found in India

3.5. Description of Basil (*Ocimum basilicum*) and Tulsi (*Ocimum sanctum*)

Ocimum is a perennial aromatic herb of the family Lamiaceae, also called Labiatae, and belongs to dicotyledons. Two hundred genera and about 3200 species are found in this family. The taxonomy of *Ocimum* is complex due to intersection hybridization and

multiplication of the species in the genus. It is a tropical plant which is cultivated and also grows as a weed. Generally, both the herbs may be propagated through seeds and sometime, they can grow through cutting stem (nodes) methods. The morphologic characteristics of both species exhibit natural variation and are separately discuss as follows:

Tulsi plant is an important symbol of Hindu religion and is worshipped and used in several religious ceremonies. *O. basilicum* is not just famous for its medicinal applications but also it is best known cooking basil.



Figure 3. Tulsi (*O. sanctum* L.) and Basil (*Ocimum basilicum* L.) in the taxonomic hierarchy under Plantae

3.5.1 Physical Features

Generally, physical features of *Ocimum* species include shape, size, color and texture of leaf, stem, seeds, inflorescence, flowers and roots.

3.5.1.1 Physical Features of *Ocimum sanctum*

Vern- *Tulsi*, Sanskrit- *Ajata*, *Maujar*, *Tulsitryum*, Eng- Holy Basil

Ocimum sanctum (synonym *Ocimum tenuiflorum*), also known as tulsi plant or holy basil is erect, much branched, sparsely hairy herb with an aromatic smell. The plant is

30- 90 cm high and the roots are tap and branched. The leaves of the plant are ovate-lanceolate with an entire margin of 2.5-6.5* 1.8-3cm. The leaves are shallowly toothed, acute or obtuse hairy with minutely dots on both surfaces. The petiole is 1-2.5cm long. The inflorescence is verticillaster. The flowers of the plant are purplish- pink which is whorled in terminal with racemes 10-15cm long. The bracts are ovate, acuminate, not exceeding the calyx. The calyx of the plant is 4mm long with upper lip broadly ovate, mucronate, recurved and with a larger lower lip. The corolla is 5-8mm long with an upper lip hairy on the back with 4 lobes and the lower lip is entire, acuminate. The seeds of the plant are globose in shape having brownish black color and fruit is carcerulus. The flowering and fruiting occur in the month from April to November.

There are two commonly known varieties of *O. sanctum*: Kali Tulsi (Krishna Tulsi) with dark purple-colored leaves and Safed Tulsi (Rama Tulsi) which has green colored leaves.

3.5.1.2 Physical Features of *Ocimum basilicum*

Vern- *Marua*, *Murya* Hindi- *Kali Tulsi*, Sanskrit- *Veruari*; Eng- Basil.

The word *basilicum* comes from the Latin word basilisk, meaning. According to Muenscher & Arthur (1978), in French, it is called as “Herbe Royale”, which suggests its positive nature. It is also called “The king of herbs” while its common name is Sweet Basil. *Ocimum basilicum* is an aromatic herb and shrub. The plant height is 50- 100 cm long. The stem is usually branched from the base with hairs. The leaves are petiolate, opposite, ovate- lanceolate, toothed or almost whole, glabrous. with a margin of 2-7*1-3cm and either green or purple. The base of the leaves is cuneate or rounded, acute or subacute, entire or toothed with dotted glands which contain volatile oil of strong smells. The inflorescence is verticillaster. The flowers of the plant are purplish or whitish- pink in color. The whorls of the flower are simple and branched racemes with a stalked bract usually shorter than calyx and is ovate and acute. The calyx of the plant is 5-8 mm long and is hairy outside with upper lip rounded and lower lip 4- dentate. The corolla of the plant is 6-8mm long and is glabrous with pubescent smell. The fruit is an achene and the nutlets of the plant are ellipsoidal in shape with a dark brown color. The leaves of *O. basilicum* are small and liniform to large, rounded and yellow-green to grey-green, red or almost black. The flowering and fruiting occur in the month from July to December.

3.6 Distribution of Basil (*Ocimum basilicum*) and Tulsi (*Ocimum sanctum*)

Ocimum is an annual plant widely found in the tropical, subtropical and temperate parts of the world. *Ocimum sanctum* is found in tropical parts of Asia and has been growing in India for over 3,000 years. It is a grassy annual plant originated from Iran, Afghanistan and India. The plant is distributed in tropical and warm temperate regions of the world including Sri Lanka, W. Asia and Australia. It is grown throughout India up to 1800m in the Himalayas, and in Andaman and Nicobar Islands.

Ocimum basilicum found in the tropical, subtropical and temperate parts of the world and has especially established itself in Ceylon, hot West Asia, Africa, Malayan and

Pacific Islands. It is also found in tropical and hot temperate regions of India and Pakistan and also cultivated in several parts of US states. *Ocimum basilicum* is originally native to India and other regions of Asia. *Ocimum basilicum* L. includes annual and perennial herbs and shrubs from tropical and subtropical regions of Asia, Africa, Central and South America. The plant is usually cultivated near home yards rarely met as an escape, road sides or waste places.

4. Climatic Conditions for Plant Growth

Ocimum sanctum develops well in a wide range of soils. Therefore, poor laterite, rich loam, saline and alkaline to moderately acidic soil is appropriate for its cultivation. It grows in partial shade and thrives in fairly high rainfall and humid conditions. It can grow upto an altitude of 900m and is tolerant to drought and frost. This plant can be propagated through seeds.

Ocimum basilicum is well represented in warmer parts up to 1800m altitude above sea level. Plant cultivation can be done in wide range of soil type; dry loam from medium to fine loam. Well drained soil helps to encourage improved vegetative growth. Long day, high temperature and high humidity are found to be conducive to plant growth. Apart from regular cultivation methods, *Ocimum* needs well drained loamy to sandy loam soils with suitable pH ranging from 4.3-9.1. Watering in the form of shower is preferred which don't let the soil get hard. Soil must be moderately fertile. It can tolerate high concentrations of copper and zinc, but is more vulnerable to cobalt and nickel.

-
-

TO ACCESS ALL THE 53 PAGES OF THIS CHAPTER,
Visit: <http://www.eolss.net/Eolss-sampleAllChapter.aspx>

Bibliography

Abd El-Azim, M.H.M., Abdelgawad, A.A.M., El-Gerby, M., Ali, S., El-Mesallamy, A.M.D., (2015b), Chemical composition and antimicrobial activity of essential oil of Egyptian *Ocimum basilicum*, *Indo. Am. J. Pharm Sci.*, 2, 837-842.

Abd El-Azim, M.H.M., Abdelgawad, A.A.M., El-Gerby, M., Ali, S., El-Mesallamy, A.M.D., (2015a), Phenolic compounds and cytotoxic activities of methanol extract of Basil (*Ocimum basilicum* L.). *J. Microb. Biochem. Technol.*, 7(4): 182-185.

Abeywardhana, K.W., Abeysinghe, D.C., Dharmadasa, R.M., Aththanayake, A.M.L. (2014), Determination of optimum maturity stage for *Ocimum sanctum* grown under different growing systems in terms of therapeutically active secondary metabolites, *World J. Agric. Res.*, 2(4), 159-162.

Aburigal, Y.A.A., Mirghani, M.E.S., Elmogtaba, E.Y., Sirible, A.A.M., Hamza, N.B. Hussein, I.H. (2017). Total phenolic content and antioxidant capacity of basil (*Ocimum basilicum* L.) leaves from different locations, *Int. Food Res. J.*, 24(24), S378-S381.

Aburjai, T.A., Mansi, K., Azzam, H., Alqudah, D.A., Alshaer, W., Abuirjei, M. (2020), Chemical composition and anticancer potential of essential oil from greenhouse-cultivated *Ocimum basilicum*

leaves. *Indian J. Pharm. Sci.*, 82(1), 179-184. [In this paper antitumor activity of essential oil of *Ocimum basilicum* was analysed].

Ademiluyi, A.O., Oyeleye, S.I., Oboh, G. (2016). Biological activities, antioxidant properties and phytoconstituents of essential oil from sweet basil (*Ocimum basilicum* L.) leaves, *Comp. Clin. Pathol.*, 25, 169-176.

Adiguzel, A., Gulluce, M., Sengul, M., Ogutcu, H., Sahin, F., Karaman, I. (2005), Antimicrobial effects of *Ocimum basilicum* (Labiatae) extract. *Turk J. Biol.*, 29, 155-160.

Agarwal, P., Nagesh, L., Murlikrishnan (2010). Evaluation of the antimicrobial activity of various concentrations of Tulsi (*Ocimum sanctum*) extract against *Streptococcus mutans*: An in vitro study, *Indian J. Dent. Res.*, 21, 357.

Agatonovic-Kustrin, S., Kustrin, E., Gegechkori, V., Morton, D.W. (2020), Anxiolytic terpenoids and aromatherapy for anxiety and depression, In: *Reviews on New Drug Targets in Age-related Disorders* (Eds. Paul C. Guest), pp. 283-296

Ahmad, K., Yusra, K., Somayya, R. (2016). Antifungal, phytotoxic and hemagglutination activity of methanolic extracts of *Ocimum basilicum*, *J. Tradit. Chin. Med.*, 36(6), 794-798.

Ahmeda, A.F., Attiac, F.A.K., Liua, Z., Lia, C., Weia, J., Kanga, W. (2019). Antioxidant activity and total phenolic content of essential oils and extracts of sweet basil (*Ocimum basilicum* L.) plants. *Food Sci. Hum. Well.*, 8, 299-305.

Akgul, A. (1989), Volatile oil composition of sweet basil (*Ocimum basilicum* L.) cultivating in Turkey, *Nahrung*, 33, 87-88.

Akoto, C.O., Acheampong, A., YawDuahBoakye, Naazo, A.A., Adomah, D.H. (2020), Anti-inflammatory, antioxidant, and anthelmintic activities of *Ocimum basilicum* (sweet basil) fruits, *J. Chem.*, Article ID 2153534.

Al-Ghamdi, A.Y., Abdalla, M.O.M., Fadlelmula, A. A. (2020), Phytochemical, total phenolic contents, and antioxidant and antimicrobial activities of *Ocimum basilicum* L. leaf extract in Al-Baha Area, Saudi Arabia. *Int. J. Adv. Res.* 8(03), 526-533.

Ali, B., Al-Wabel, N.A., Shams, S., Ahamad, A., Khan, S.A., Anwar, F. (2015), Essential oils used in aromatherapy: A systemic review. *Asian Pac. J. Trop. Biomed.*, 5(8), 589-598. [This review has shown direct influence on the application parts of the essential oils and explore about the major plants which oil utilized in aromatherapy].

Ali, H., Dixit, S., (2012), In vitro antimicrobial activity of flavonoids of *Ocimum sanctum* with synergistic effect of their combined form, *Asian Pac. J. Trop. Dis.*, S396-S398.

Almatroodi, S.A., Alsahli, M.A., Almatroodi, A. Rahmani, A.H. (2020), *Ocimum sanctum*: Role in diseases management through modulating various biological activities, *Pharmacogn. J.*, 12(5), 1198-1205.

Amjad, K. (2013), Antimicrobial activity of ethanolic extract of *Ocimum basilicum* leaf from Saudi Arabia. *Biotechnology*, 12(1), 61-64.

Anand, A.K., Manindra, M., Haider, S.Z., Sharma, A. (2011), Essential oil composition and antimicrobial activity of three *Ocimum* species from Uttarakhand (India), *Int. J. Pharm. Sci.*, 3, 223-225.

Anand, T., Sundararajan, M., Anbukkarasi, M., Thomas, P.A., Geraldine P.A. (2019), Methanolic extract of *Ocimum basilicum* exhibits antioxidant effects and prevents selenite-induced cataract formation in cultured lenses of wistar rats. *Pharmacogn. J.* 11(3), 496-504.

Andola, H.C., Purohit, V.K., Chalum, K.A., Rawat, M.S.M. (2018). Essentials oil composition of industrially important *Ocimum basilicum* cultivated by farmers of Uttarakhand. *J. Herb Med.*, 2(2), 1-4.

Aoshima, H., Hamamoto, K. (1999). Potentiation of GABA receptors expressed in *Xenopus* oocytes by perfume and phytoncide. *Biosci. Biotechnol. Biochem.*, 63, 743-748.

Aprotosoae, A.C., Hancianu, M., Costache, I.I., Miron, A. (2014), Linalool: A review on a key odorant molecule with valuable biological properties. *Journal of Flavour and Fragrance*, 29, 193-219.

- Arimura, G., Ozawa, R., Shimoda, T., Nishioka, T., Boland, W., Takabayashi, J. (2000), Herbivory-induced volatiles elicit defense genes in lima bean leaves. *Nature*, 406, 512-515.
- Asha, M.K., Prashanth, D., Murali, B., Padmaja, R., Amit, A. (2001), Anthelmintic activity of essential oil of *Ocimum sanctum* and eugenol, *Fitoterapia*, 72(6), 669-670. [Potent anthelmintic activity of essential oil of *Ocimum sanctum* and eugenol in the *Caenorhabditis elegans* model has been investigated in this paper].
- Aslan, I., Ozbek, H., Calmasur, O., Sahin, F. (2004), Toxicity of essential oil vapours to two greenhouse pests, *Tetranychus urticae* Koch and *Bemisia tabaci* Genn. *Ind. Crops Prod.*, 19(2), 167-173.
- Atanda, O.O., Akpan, I., Oluwafemi, F. (2007), The potential of some spice essential oils in the control of *A. parasiticus* CFR 223 and aflatoxin production. *Food Control*, 18(5), 601-607.
- Awasthi, P.K., Dixit, S.C. (2007), Chemical compositions of *Ocimum sanctum* Shyama and *Ocimum sanctum* Rama oils from the plains of Northern India, *J. Essent. Oil Bear. Pl.*, 10, 292-296.
- Aye Aye, Jeon, Y.D., Lee, J.H., Bang, K.S., Jin, J.S. (2019). Anti-inflammatory activity of ethanol extract of leaf and leaf callus of basil (*Ocimum basilicum* L.) on RAW 264.7 macrophage cells. *Orient. Pharm. Exp. Med.*, 19, 217-226.
- Azmi, K.A., Ahmed, W., Siddiqui, M.K. (1999), Aromatic drugs in Unani medicine with special reference to Kitabul-Mia-Lil-Masihi. *Bull. Indian Inst. Hist. Med/ Hyderabad*, 29(2), 103-12. [The reported data described the importances of aromatic plants used as a herbal drug in Unani medicine systems with special references of a Unani book].
- Azuma, H., Thien, L.B., Kawano, S. (1999), Floral scents, leaf volatiles and thermogenic flowers in Magnoliaceae. *Plant Species Biol.*, 14, 121-127.
- Bakkali, F., Averbeck, S., Averbeck, D., Idaomar, M. (2008), Biological effects of essential oils: a review. *Food Chem. Toxicol.*, 46, 446-475. [This review explores the available information regarding biological potential of essential oils as an antioxidant, antimicrobial, antidiabetic and many more].
- Bakr, R.F., Fattah, H.M., Salim, N.M., Atiya, N.H. (2010), Insecticidal activity of four volatile oils on two museum insects pests. *Egypt. Acad. J. Biol. Sci.*, 2, 57-66.
- Balaji, R., Prakash, G., Devi, P.S. Aravinthan K. (2011). Antioxidant activity of methanol extract of *Ocimum tenuiflorum* (dried leaf and stem). *Int. J. Pharm. Res. Dev.*, 3, 20-27.
- Balakrishna, V., Pamu, S., Pawar, D. (2017), Evaluation of anti-pyretic activity of *Ocimum sanctum* L. using brewer's yeast induced pyrexia in albino rats. *J. Innovations Pharm Sci.*, 1(2), 55-58.
- Baliga, M.S., Rao, S., Rai, M.P., D'souza P. (2016), Radio protective effects of the ayurvedic medicinal plant *Ocimum sanctum* Linn. (Holy Basil): A memoir. *J. Cancer Res. Ther.* 12(1), 20-27.
- Banerjee, S., Prashar, R., Kumar, A., Rao A.R. (1996), Modulatory influence of alcoholic extract of *Ocimum* leaves on carcinogen induced metabolizing enzyme activities and reduced glutathione levels in mouse. *Nutr. Cancer.*, 25(2), 205-217.
- Bano, N., Ahmed, A., Tanveer, M., Khan, G.M., Ansari, M.T. (2017), Pharmacological evaluation of *Ocimum sanctum*. *J. Bioequiv. Availab.*, 9:3. [All pharmaceutical potential of *Ocimum sanctum* has deeply discussed here].
- Bansavatar, C.S., Kurup, R., Ansari, A.A. (2015), Antimicrobial properties of *Ocimum sanctum* and *Calotropis gigantea* leaves. *Br. Microbiol. Res. J.*, 8(31), 532-539.
- Bansod, S., Rai, M. (2008), Antifungal Activity of essential oils from Indian medicinal plants against human pathogenic *Aspergillus fumigatus* and *A. niger*. *World J. Medical Sci.*, 3(2), 81.
- Barcelos, R.C., Jham, G.N., Onkar, D., Fernanda, A., Vania, M., Valente, M.M. (2013), Identification and quantification of the major fungi toxic components of the Brazilian basil (*Ocimum basilicum* L.) essential oil, *J. Food Res.*, 2, 5.
- Behbahani, M. (2014), Evaluation of *in vitro* anticancer activity of *Ocimum basilicum*, *Alhagi maurorum*, *Calendula officinalis* and their parasite *Cuscuta campestris*, *PLoS ONE* 9(12), e116049.
- Benedec, D., Parvu, A.E., Oniga, I., Toiu, A., Tiperciuc, B. (2007). Effects of *Ocimum basilicum* L. extract on experimental acute inflammation. *Rev. Med. Chir. So.c Med. Nat. Iasi.*, 111, 1065- 1069.

- Benedec, D., Vlase, L., Hanganu, D., Oniga, I. (2012). Antioxidant potential and polyphenolic content of Romanian *Ocimum basilicum*. *Dig. J. Nanomater. Bios.*, 7(3), 1263-1270.
- Bhatt, S., Tewari, G., Pande C., Rana L. (2018), Impact of drying methods on essential oil composition of *Ocimum americanum* L. from Kumaun Himalayas, *J. Essent. Oil Bear. Pl.*, 21, 1385-1396.
- Bhattacharya, A., Aggarwal, A., Sharma, N., Cheema, J. (2014), Evaluation of some anti-oxidative constituents of three species of *Ocimum*, *Int. J. Life Sci.*, 8, 14-17.
- Bhattacharyya, D., Sur, T.K., Jana, U., Debnath, P.K. (2008). Controlled programmed trial of *Ocimum sanctum* leaf on generalized anxiety disorders. *Nepal Med. Coll. J.*, 10, 176 179.
- Black, R.E., Cousens, S., Johnson, H.L. (2010), Global, regional, and national causes of child mortality in 2008: A systematic analysis. *Lancet*, 375, 1969-1987.
- Boateng, J., Catanzano, O. (2015). Advanced therapeutic dressing for effective wound healing-review, *J. Pharm. Sci.*, 104(11), 3653-3680.
- Bob, H. (2002), Methyl Eugenol-the current bete noire of aromatherapy, *Int. J. Aromather.*, 12, 193-201.
- Borah, R., Biswas, S.P. (2018), Tulsi (*Ocimum sanctum*), excellent source of phytochemicals, *Int. J. Environ. Agric. Biotech.*, 3(5), 1732-1738.
- Borges, S.R., Ortiz, B.L.S., Pereira, A.C.M., Keita, H., Carvalho, J.C.T. (2019). *Rosmarinus officinalis* essential oil: a review of its phytochemistry, anti-inflammatory activity, and mechanisms of action involved. *J. Ethnopharmacol.*, 229, 29-45.
- Boukhatem, M.N., Setzer, W.N. (2020). Aromatic herbs, medicinal plant-derived essential oils, and phytochemical extracts as potential therapies for coronaviruses: Future Perspectives. *Plants*, 9(6), 800.
- Bouvier-Brown, N., Goldstein, A., Worton, D., Matross, D., Gilman, J., Kuster, W., Welsh-Bon, D., Warneke, C., Gouw, J., Cahill, T., Holzinger, R. (2008), Methyl chavicol: characterization of its biogenic emission rate, abundance, and oxidation products in the atmosphere. *Atmos. Chem. Phys. Discuss.*, 8, 19707-19741.
- Bouwmeester, H., Dekkers, S., Noordam, M.Y., Hagens, W.I., Bulder, A.S., De Heer, C., Ten Voorde, S.E., Wijnhoven, S.W., Marvin, H.J., Sips, A.J. (2009). Review of health safety aspects of nanotechnologies in food production. *Regul. Toxicol. Pharmacol.*, 53, 52-62
- Buchbauer, G., Jirovetz, L., Jager, W., Dietrich, H., Plank, C. (1991) Aromatherapy: evidence for sedative effects of the essential oil of lavender after inhalation. *Z. Naturforsch. C.*, 46(11-12), 1067-1072.
- Buddhadev, S.G., Buddhadev, S.S., Mehta, N.D. (2014). A review article on *Ocimum sanctum* L. *Punarnav: Int. Peer Rev. Ayurved. J.*, 2(2), 1-6.
- Bunrathep, S., Palanuvez, C., Ruangrunsi, N. (2007), Chemical composition and antioxidative activities of essential oils from four *Ocimum* species endemic to Thailand, *J. Health Res.*, 21, 201-206.
- Burt, S. (2004). Essential oils: Their antibacterial properties and potential applications in foods - A review. *J. Agric. Food Chem.*, 94, 223-253. [This study focused on the application of essential oils in food as preservative owing to its antibacterial effects].
- Cha, J.H., Kim, M.J., Kim, H.S., Kim, Y.I. (2010). Effects of aromatherapy in lending oil of basil, lavender, rosemary and rose on headache, anxiety and serum cortisol level in the middle-aged women. *J. Korean Biol. Nurs. Sci.*, 12(3), 133-139.
- Chandrappa, P.M., Dupper, A., Tripathi, P., Arroju, R., Sharma, P. (2015) Antimicrobial activity of herbal medicines (tulsi extract, neem extract) and chlorhexidine against *Enterococcus faecalis* in Endodontics: An *in vitro* study. *J. Int. Soc. Prev. Commun. Dent.*, 5, S89-S92.
- Chang, C.L., Cho, I.K., Li, Q.X. (2009b). Insecticidal activity of basil oil, *trans*-anethole, estragole, and linalool to adult fruit flies of *Ceratitis capitata*, *Bactrocera dorsalis*, and *Bactrocera cucurbitae*. *J. Econ. Entomol.*, 102, 203-209.
- Chang, X., Alderson, P.G., Wright, C.J., (2009a), Variation in the essential oils in different leaves of basil (*Ocimum basilicum* L.) at day time, *Open Hort. J.*, 2, 13-16.

- Chaudhary, A., Sharma, S., Mittal, A., Gupta, S., Dua, A. (2020), Phytochemical and antioxidant profiling of *Ocimum sanctum*. *J. Food Sci. Technol.* 57, 3852-3863.
- Chiang, L.C., Ng, L.T., Cheng, P.W., Chiang, W., Lin, C.C. (2005), Antiviral activities of extracts and selected pure constituents of *Ocimum basilicum*, *Clin. Exp. Pharmacol. Physiol.*, 32(10), 811-816.
- Chil, N. I., Escalona A.J.C, Berenguer, R.C.A, Mendonça, P.M, Mateo –Perez, K., Dutok, S.C. M., Cortinhas, L.B., Silva, C.F, Carvalho., M.G, Queiroz, M.M.C. (2017), Chemical Composition and Toxicity of *Ocimum sanctum* L. Var. *Cubensis*. Essential oil up-growing in the eastern of Cuba, *Int. J. Pharmacogn. Phytochem. Res.*, 9(7), 1021-1028
- Chowdhury, I. I., Rahman, Md. A., Hashem, M. A., Bhuiyan, M. M. H., Hajjar, D., Alelwani, W., Makki, A.A., Haque, Md. A., Tangpong, J., Bakhtiar, M. T. B. (2020). Supplements of an aqueous combination of *Justicia adhatoda* and *Ocimum tenuiflorum* boost antioxidative effects and impede hyperlipidemia. *Animal Model. Exp. Med.*, 29(3), 140-151.
- Christman, S. 2010. *Ocimum basilicum*. *Floridata* Tallahassee, FL, available at: <http://www.floridata.com/ref/o/ocim-bas.cfm>.
- Clark, G.S. (1988). A profile: An aroma chemical - *Linalool*. *Perfum. Flavor.*, 13(8-9), 49-54. [The proposed chapter discussed detailed study on the compound linalool which was major component of *Ocimum* extract as well as essential oil].
- Coats, J., Schultz, G., Peterson, C. (2003), In: *Botanical products as repellents against mosquitoes and cockroaches.*, 226th American Chemical Society National Meeting, New York, American Chemical Society, Washington, D.C. New York.
- Da Silva, J.K.R., Figueiredo, P.L.B., Byler K.G. Setzer, W.N. (2020). Essential oils as antiviral agents, potential of essential oils to treat SARS-CoV-2 Infection: An in-silico investigation. *Int. J. Mol. Sci.*, 21, 3426.
- Dahanukar, S.A., Kulkarni, R.A., Rege N.N. (2000), Pharmacology of medicinal plants and natural products, *Indian J. Pharmacol.*, 32, 81-118. [Active agents present in the natural medicinal plants have been discussed in this paper].
- Dambolena, J.S., Zunino, M.P., López, A.G., Rubinstein, H.R., Zygadlo, J.A., Mwangi, J.W., Thoithi, G.N., Kibwage, I.O., Mwalukumbi, J.M., Kariuki, S.T. (2010), Essential oils composition of *Ocimum basilicum* L. and *Ocimum gratissimum* L. from Kenya and their inhibitory effects on growth and fumonisin production by *Fusarium verticillioides*, *Innov. Food Sci. Emerg. Technol.*, 11, 410-414.
- Das, J., Buragohain, B., Srivastava, R.B. (2010). In vitro evaluation of *Ocimum sanctum* leaf extract against dermatophytes and opportunistic fungi. *Asian J. Microbiol., Biotechnol. Environ. Sci.*, 12, 789-792.
- Das, M.K., Mandal, M., Mandal, S. (2017). Broad antibacterial spectrum and high performance liquid chromatography profiles of *Ocimum sanctum* leaf extract. *Acta Sci. pharm.Sci.* 1(6): 02-07.
- De Lima Gondim, F., dos Santos, G.R., do Nascimento, I.F.M.G., Serra, D.S., Cavalcante, F.S.A. (2018). Beneficial effects of eucalyptol in the pathophysiological changes of the respiratory system: a proposal for alternative pharmacological therapy for individuals with COPD. *European Journal of Medicinal Plants*, 25(1), 1-10.
- De Sousa, D.P., Hocayen, P.D.A.S., Andrade, L.N., Andreatini, R. (2015), A systematic review of the anxiolytic-like effects of essential oils in animal models. *Molecule*, 20, 18620-18660.
- Devi, P.U. (2001). Radioprotective, anticarcinogenic and antioxidant properties of the Indian holy basil, *Ocimum sanctum* (Tulasi), *Indian J. Exp. Biol.*, 39, 185-190.
- Dhif, W., Bellili, S., Jazi, S., Bahloul, N., Mnif, W. (2016), Essential oils chemical characterization and investigation of some biological activities: A critical review. *Medicines*, 3(4), 22-25.
- Dohare, S.L., Shuaib, S., Ahmad, M.I., Naquv, K.J. (2012). Chemical composition of volatile oil of *Ocimum sanctum* linn., *Int. J. Biomed. Res.*, 3(2), 129-131.

Dötterl, S., Burkhardt, D., Weibecker, B., Jürgens, A., Schütz, S., Mosandl, A. (2006), Linalool and lilac aldehyde/alcohol in flower scents electrophysiological detection of lilac aldehyde stereoisomers by a moth. *J. Chromatogr. A.*, 1113, 231-238.

Dreher, F., Maibach, H.I. (2001), Protective effects of topical antioxidants in humans, In *Oxidants and Antioxidants in Cutaneous Biology* (Eds. J.J. Thiele, P. Elsner), Basel: Karger 15-164 pp. [This chapter summarised the responsibilities and photoprotective effects of ‘natural’ and synthetic antioxidants as skin care agents].

Dris, D., Tine-Djebbar, F., Bouabida, H., Soltani, N. (2017), Chemical composition and activity of an *Ocimum basilicum* essential oil on *Culex pipiens* larvae: Toxicology, biometrical and biochemical aspects. *S. Afr. J. Bot.*, 113, 362-369.

Drulis-Kawa, Z., Dorotkiewicz-Jach, A. (2010). Liposomes as delivery systems for antibiotics, *Int. J. Pharm.*, 387(1), 187-198. Eftekhari, N., Moghimi, A., Roshan, M.N., Saadat, S., Boskabady, M.H. (2019), Immunomodulatory and anti-inflammatory effects of hydro-ethanolic extract of *Ocimum basilicum* leaves and its effect on lung pathological changes in an ovalbumin-induced rat model of asthma. *BMC Complement. Altern. Med.*, 19, 349.

Elisabetsky, E., Brum, L., Souza, D. (1995). Anticonvulsant properties of linalool in glutamate-related seizure models. *Phytomedicine*, 6, 107-113.

Fakhar, Z., Mostofi, Y., Zamani, Z. (2014), Application of *Ocimum basilicum* essential oil as vapor on postharvest storage of plum fruit cv. ‘Golden Drop’. *Not. Sci. Biol.*, 6(4), 454-459.

Fang, J.Y., Leu, Y., Hwang, T., Cheng, H. (2004). Essential oils from sweet basil (*Ocimum basilicum*) as novel enhancers to accelerate transdermal drug delivery. *Biol. Pharma. Bull.*, 27, 1819-1825.

Fathiazad, F., Matlobi, A., Khorrami, A., Hamedeyazdan, S., Soraya, H., Hammami, M., Maleki-Dizaji, N., Garjani, A. (2012), Phytochemical screening and evaluation of cardioprotective activity of ethanolic extract of *Ocimum basilicum* L. (basil) against isoproterenol induced myocardial infarction in rats. *J. Pharm. Sci.*, 20(1), 87.

FEMA (1997), FEMA Database: Linalool (FEMA No. 2635), Washington, DC, Flavor and Extract Manufacturers’ Association, pp. 53.

Fowler, M.W., Scragg, A.H. (1988), Natural products from higher plants and plant cell culture, In: *Plant cell Biotechnology* (Eds. M.S.S. Pais, F. Mavituna, J.M. Novais). NATO ASI Series. Berlin: Springer-Verlag; 18: 165-77pp.

Gadiyar, A., Ankola, A.V., Rajpurohit, L.S. (2017). Evaluation of the antimicrobial activity of *Ocimum sanctum* L. (Tulsi) extract against *Streptococcus Mutans* and *Lactobacillus acidophilus* - An in vitro study. *International Journal of Health Sciences and Research*, 7(4), 224.

Gaio, I., Saggiolato, A.G., Treiche, H., Cichoski, A.J., Astolfi, V., Cardoso, R.I., Toniazzi, G., Valduga, E., Paroul, N., Cansian, R.L. (2015), Antibacterial activity of basil essential oil (*Ocimum basilicum* L.) in Italian-type sausage, *J. Verbr. Lebensm. J. Consum., Prot. Food S.* 10(4), 323-329.

Gaofeng, Y., Wahlqvist, M.L., He, G., Yang, M., Li, D. (2006). Natural products and anti-inflammatory activity. *Asia Pac. J. Clin. Nutr.*, 15(2), 143-152. [This review summarized anti-inflammatory efficiency of the components of natural products].

Gaur. R.D. (1999), Flora of District Grahwal North west Himalayas, *Transmedia*: Srinagar, Garhwal.

Gavanji, S., Mohabatkar, H., Baghshahi, H., Zarrabi, A. (2014), Bioinformatics prediction of interaction silver nanoparticles on the disulfide bonds of HIV-1 Gp120 protein. *International Journal of Scientific Research and Knowledge*, 2, 67-74.

Gengan, R.M., Anand, K., Phulukdaree, A., Chuturgoon, A. (2013), A549 lung cell line activity of biosynthesized silver nanoparticles using *Albizia adianthifolia* leaf, *Colloids Surf. B.*, 105, 87-91.

Ghoke, S.S., Sood, R., Kumar, N., Pateriya, A. K., Bhatia, S., Mishra, A., Dixit, R., Singh, V.K., Desai, D.N., Kulkarni, D.D., Dimri, U., Singh, V.P. (2018), Evaluation of antiviral activity of *Ocimum sanctum* and *Acacia arabica* leaves extracts against H9N2 virus using embryonated chicken egg model, *BMC Complement. Altern. Med.*, 18, 174.

- Ghosh V., Mukherjee A., Chandrasekaran N. (2013), Ultrasonic emulsification of food-grade nanoemulsion formulation and evaluation of its bactericidal activity, *Ultrasonics Sonochemistry*, 20, 338-344.
- Gkinis, G., Tzakou, O., Iliopoulou, D., Roussis, V. (2003), Chemical composition and biological activity of Nepeta parnassica oils and isolated nepetalactones, *J. Biosci.*, 58(9/10), 681-686.
- Godhwani, S., Godhwani, J.L., Was. D.S. (1988), O. sanctum- A preliminary study evaluating its immunoregulatory prole in albino rats, *J. Ethnopharmacol.*, 24, 193-198.
- Goel, A., Kumar, S., Singh, D.K., Bhatia, A.K. (2010), Wound healing potential of Ocimum sanctum Linn. with induction of tumor necrosis factor-alpha, *Indian J. Exp. Biol.*, 48, 402-406.
- Goyal, P., Kaushik, P. (2011), In vitro evaluation of antibacterial activity of various crude leaf extracts of Indian sacred plant, *Ocimum sanctum* L. Br. *Microbiol. Res. J.*, 1, 70-8.
- Grabowska, K., Janeczko, Z. (2013), Olejki eteryczne w preparatach farmaceutycznych. *Aromaterapia*, 4(74), 16-50.
- Gradinariu, V., Cioanca, O., Hritcu, L., Trifan, A., Gille, E., Hancianu, M. (2015), Comparative efficacy of *Ocimum sanctum* L. and *Ocimum basilicum* L. essential oils against amyloid beta (1-42)- induced anxiety and depression in laboratory rats. *Phytochem. Rev.*, 14, 567-575.
- Gucwa, K., Milewski, S., Dymerski, T., Szveda, P. (2018). Investigation of the antifungal activity and mode of action of *Thymus vulgaris*, *Citrus limonum*, *Pelargonium graveolens*, *Cinnamomum cassia*, *Ocimum basilicum*, and *Eugenia caryophyllus* essential oils, *Molecules*, 23(5), 1116.
- Guez, C.M., De-Souza, R.O., Fischer, P., Leao-De, M.M.F., Duaetr, J.A., Boligon, A.A., Athayde, M.L., Zuravski, L., Oliveira-DeSouza, L.F., Machado, M.M. (2017). Evaluation of basil extract (*Ocimum basilicum* L.) on oxidative, anti-genotoxic and anti-inflammatory effects in human leukocytes cell cultures exposed to challenging agents. *Braz. J. Pharm. Sci.*, 53(1), 1-12. e15098.
- Guimarães, A.C., Meireles, L., Lemos, M.F. Guimarães, M.C.C., Endringer, D.C., Fronza, M., Scherer, R. (2019), Antibacterial activity of terpenes and terpenoids present in essential oils. *Molecules*, 24, 2471.
- Gulçin, I. Elmastas M., Aboul-Enein, H.Y. (2007), Determination of antioxidant and radical scavenging activity of basil (*Ocimum basilicum* L. Family Lamiaceae) assayed by different methodologies. *Phytother. Res.*, 21, 354-361
- Guo, D., Zhu, L., Huang, Z., Zhou, H., Ge, Y., Ma, W., Wu, J., Zhang, X., Zhou, X., Zhang, Y., Zhao, Y., Gu, N. (2013), Anti-leukemia activity of PVP-coated silver nanoparticles via generation of reactive oxygen species and release of silver ions, *Biomaterials*, 34, 7884-7894.
- Gupta, S.K., Prakash, J., Srivastav, S. (2002), Validation of claim of Tulsi, *Ocimum sanctum* Linn as a medicinal plant, *J. Expt. Biol.*, 40, 765-773.
- Gupta, P., Yadav, D.K., Siripurapu, K.B., Palit, G., Maurya, R. (2007), Constituents of *Ocimum sanctum* with antistress activity, *J. Nat. Prod.*, 70(9), 1410-1416.
- Habibu, T., Abubakar, M., Ladidi, I.Z., Adegbenro, A.P., Carrol, L.D., Jibril, A.A. (2017), *Current Status of Antidiabetic Plants in the Lamiaceae family*, Editions Universitaires Europeennes.
- Hakkim, F.L., Shankar, C.G., Girua, S. (2007). Chemical composition and antioxidant property of holy basil (*Ocimum sanctum* L.) leaves, stems, and inflorescence and their *in vitro* callus cultures. *J. Agr. Food Chem.*, 55, 9109-9117.
- Han, H.D., Cho, Y.J., Cho, S.K., Byeon, Y., Jeon, H.N., Kim, H.Y., Kim, B.G., Bae, D.S., Lopez-Berestein, G., Sood, A.K., Shin, B.C., Park, Y.M., Lee, J.W. (2016), Linalool-incorporated nanoparticles as a novel anticancer agent for epithelial ovarian carcinoma. *Mol. Cancer Ther.*, 15(4), 618-627.
- Han, X., Beaumont, C., Stevens, N. (2017), Chemical composition analysis and in vitro biological activities of ten essential oils in human skin cells, *Biochim. Open*, 5, 1-7.
- Hanif, M.A., Al-Maskari, M.Y., Al-Maskari, A., Al-Shukaili, A., Al-Maskari, A.Y., Al-Sabahi, J.N. (2011), Essential oil composition, antimicrobial and antioxidant activities of unexplored Omani basil, *J. Med. Plant. Res.*, 5, 751-758.

Hannan, J.M.A., Das, B.K., Uddin, A., Bhattacharjee, R., Das, B., Chowdury, H.S., Mosaddek, A.S.M. (2011), Analgesic and anti-inflammatory effects of *Ocimum sanctum* (linn) in laboratory animals, *International Journal of Pharmaceutical Sciences and Research*, 2(8), 2121-2125.

Harnafi, H., Aziz, M., Amrani, S. (2009), Sweet basil (*Ocimum basilicum* L.) improves lipid metabolism in hypercholesterolemic rats. *Eur. E J. Clin. Nutr. Metab.*, 4, 181-186.

Hopp, R., Mori, K. (1993), Recent Developments in Flavour and Fragrance Chemistry, Proceedings of the 3rd International Haarmann and Reimer Symposium, VCH Verlagsgesellschaft Germany, pp: 123-128. <https://en.wikipedia.org/wiki/Ocimum>

https://gardener.fandom.com/wiki/Sweet_basil

Huang, H.C., Ho, Y.C., Lim, J.M., Chang, T.Y., Ho, C.L., Chang, T.M. (2015). Investigation of the anti-melanogenic and antioxidant characteristics of *Eucalyptus camaldulensis* flower essential oil and determination of its chemical composition. *Int. J. Mol. Sci.*, 16, 10470-10490.

Hussain, A.I., Anwar, F., Sherazi, S.T.H., Przybylski, R. (2008), Chemical composition, antioxidant and antimicrobial activities of basil (*Ocimum basilicum*) essential oils depends on seasonal variations, *Food Chem.*, 108, 986-995.

Hussain, A.I., Chatha, S.A.S., Kamal, G.M., Ali, M.A., Hanif, M.A., Lazhari, M.I. (2016), Chemical composition and biological activities of essential oil and extracts from *Ocimum sanctum*. *Int. J. Food Prop.*, 20(7), 1969-1981.

Hussain, A.I., Chatha, S.A.S., Kamal, G.M., Ali, M.A., Hanif, M.A., Lazhari, M.I. (2017), Chemical composition and biological activities of essential oil and extracts from *Ocimum sanctum*, *Int. J. Food Prop.*, 20(7), 1569-1581.

Iqbal, Z., Akhtar, M., Sabri, M. U., Altaf, A. (2020), Chemical composition of *Ocimum sanctum* essential oil by GC-MS analysis, *Nat. Prod. Chem.* 8(6), 1-5.

Ismail, M. (2006), Central properties and chemical composition of *Ocimum basilicum* essential oil, *Pharm. Biol.*, 44, 619-626.

Jabir, M., Sahib, U.I., Taqi, Z., Taha, A., Sulaiman, G., Albukhaty, S., Al-Shammari, A., Alwahibi, M., Soliman, D., Dewir, Y.H., Rizwana, H. (2020), Linalool-loaded glutathione-modified gold nanoparticles conjugated with CALNN peptide as apoptosis inducer and NF-KB translocation inhibitor in SKOV-3 cell line. *Int. J. Nanomed.*, 15, 9025-9047.

Jaggi, R.K., Madaan, R., Singh, B. (2003). Anticonvulsant potential of holy basil, *Ocimum sanctum* Linn., and its cultures. *Indian J. Exp. Biol.*, 41, 1329-1333.

Jalilzadeh-Amin, G., Maham, M. (2015), The application of 1,8- cineole, a terpenoid oxide present in medicinal plants, inhibits castor oil-induced diarrhea in rats. *Pharm. Biol.*, 53(4), 594-599.

Jayanti, I., Jalaluddin, M., Avijeeta, A., Ramanna, P.K., Rai, P.M., Nair, R.A. (2018), *In vitro* antimicrobial activity of *Ocimum sanctum* (Tulsi) extract on *Aggregatibacter actinomycetemcomitans* and *Porphyromonas gingivalis*. *J. Contemp. Dent. Pract.*, 19(4), 415-419.

Jayasinghe, C., Gotoh, N., Aoki, T., Wada, S. (2003), Phenolics Composition and antioxidant activity of sweet basil (*Ocimum basilicum* L.), *J. Agric. Food Chem.*, 51, 4442-4449.

Jirovetz, L., Buchbauer, G., Stoyanova, A., Balinova, A. (2001). Analysis, chemotype and quality control of the essential oil of a new cultivated basil (*Ocimum basilicum* L.) plant from Bulgaria. *Sci. Pharm.*, 69(1), 85-89.

Joshi R.K. (2014). Chemical composition and antimicrobial activity of the essential oil of *Ocimum basilicum* L. (sweet basil) from Western Ghats of North West Karnataka, India. *Anc. Sci. Life*, 33(3), 151.

Joshi, R.K. (2013), Chemical composition, *in vitro* antimicrobial and antioxidant activities of the essential oils of *Ocimum gratissimum*, *O. sanctum* and their major constituents, *Indian J. Pharm. Sci.*, 75(4): 457-462.

Juergens, L.J., Worth, H., Juergens, U.R. (2020). New Perspectives for mucolytic, anti-inflammatory and adjunctive therapy with 1,8-cineole in COPD and asthma: Review on the new therapeutic approach. *Adv. Ther.*, 37(5), 1737-1753.

- Juntachote, T., Bergofer, E. (2005), Antioxidative properties and stability of ethanolic extracts of Holy basil and Galangal. *Food Chem.*, 92, 193-202.
- Kadan, S., Saad, B., Sasson, Y., Zaid, H. (2016). In vitro evaluation of anti-diabetic activity and cytotoxicity of chemically analysed *Ocimum basilicum* extracts. *Food Chem.* 196, 1066-1074.
- Kamatou, G.P., Viljoen, A.M. (2008), Linalool-a review of a biologically active compound of commercial importance. *Nat. Prod. Commun.*, 3, 1183-1192.
- Kamyab, A.A., Eshraghian, A. (2013). Anti-Inflammatory, gastrointestinal and hepatoprotective effects of *Ocimum sanctum* Linn: an ancient remedy with new application. *Inflamm. Allergy Drug Targets*, 12, 378-384.
- Kang, Z.W., Liu, F.H., Zhang, Z.F., Tian, H.G., Liu, T.X. (2018). Volatile β -ocimene can regulate developmental performance of Peach Aphid *Myzus persicae* through activation of defense responses in Chinese cabbage *Brassica pekinensis*. *Front. Plant Sci.*, 9, 1-12.
- Karthikeyan, K., Gunasekaran, P., Ramamurthy, N., Govindasamy, S. (1999), Anticancer activity of *Ocimum sanctum*, *Pharm. Biol.*, 37(4), 285-290.
- Kasali, A.A., Eshilokun, A.O., Adeola, S., Winterhalter, P., Knapp, H., Bonnlander, B., Koenig, W.A. (2005), Volatile oil composition of new chemotype of *Ocimum basilicum* L. from Nigeria, *Flavour Frag. J.*, 20, 45- 47.
- Kath, R.K. Gupta, R.K., (2006). Antioxidant activity of hydroalcoholic extract of *Ocimum sanctum* in animal models of peptic ulcer. *Indian J. physiol. Pharmacol.*, 50(4), 391-396.
- Kathirvel, P., Ravi, S. (2012). Chemical composition of the essential oil from basil (*Ocimum basilicum* Linn.) and its *in vitro* cytotoxicity against HeLa and Hep-2 human cancer cell lines and NIH 3T3 mouse embryonic fibroblasts. *Nat. Prod. Res.*, 26(12), 1112-1118.
- Kaurinovic, B., Popovic, M., Vlasisavljevic, S., Trivic, S. (2011), Antioxidant capacity of *Ocimum basilicum* L. and *Origanum vulgare* L. extracts. *Molecules*, 16, 7401-7414.
- Kavitha, S., John, F., Indira, M. (2015), Amelioration of inflammation by phenolic rich methanolic extract of *Ocimum sanctum* leaves in isoproterenol induced myocardial infraction. *Indian J. Exp. Bot.*, 53(10), 632-640.
- Kehrl, W., Sonnemann, U., Dethlefsen, U. (2004) Therapy for acute nonpurulent rhinosinusitis with cineole: results of a double-blind, randomized, placebo-controlled trial. *Laryngoscope*, 114, 738-742.
- Keita, S.M., Vincent, C., Schmit, J.P., Belanger, A. (2000), Essential oil composition of *Ocimum basilicum* L., *Ocimum gratissimum* L. and *Ocimum suave* L. in the Republic of Guinea, *Flavour Frag. J.*, 15, 339-341.
- Kelm, M.A., Nair, M.G., Strasburg, G.M., DeWitt, D.L. (2000), Antioxidant and cyclooxygenase inhibitory phenolic compounds from *Ocimum sanctum* Linn. *Phytomedicine*, 7, 7-13.
- Khan, A., Ahmad, A., Manzoor, N., Khan, L.A. (2010a), Antifungal activities of *Ocimum sanctum* essential oil and its lead molecules, *Nat. Prod. Commun.*, 5(2), 345-349.
- Khan, A., Ahmed, A., Akhtar, F., Yousuf, S., Xess, I., Khan, L.A., Manzoor, N. (2010c), *Ocimum sanctum* essential oil and its active principles exert their antifungal activity by disrupting ergosterol biosynthesis and membrane integrity. *Res. Microbiol.*, 161(10), 816-823.
- Khan, M.R., Islam, M.A., Hossain, M.S., Asadujjaman, M., Wahed, M.I., Rahman, B.M., Anisuzzaman, A.S.M., Shaheen, S.M., Ahmed, M. (2010b), Antidiabetic effects of the different fractions of ethanolic extracts of *Ocimum sanctum* in normal and alloxan induced diabetic rats. *J Sci Res*, 2(1), 158-168.
- Khatri, L.M., Nasir, M.K., Saleem, R., Noor, F., Saleem, M.K., Noor, R.F. (1995), Evaluation of Pakistani sweet basil oil for commercial exploitation. *Pak. J. Sci. Ind. Res.*, 38, 281-282.
- Khelifa, L.H., Brada, M., Brahmi, F., Achour, D., Fauconnier, M.L., Lognay G. (2012), Chemical composition and antioxidant activity of essential oil of *Ocimum basilicum* leaves from the northern region of Algeria. *J. Herb. Med.*, 1(2), 53-58.

- Kieltyka-Dadasiewicz, A., Gorzel, M. (2014), Alternative therapies. Aromatherapy-raw materials and treatments, *European Journal of Medical Technologies*, 1(2), 72-79.
- Kim, H.J., Chen, F., Wang X., Rajapakse, N.C. (2006), Effect of methyl jasmonate on secondary metabolites of sweet basil (*Ocimum basilicum* L.). *J. Agric. Food Chem.*, 54(6), 2327-2332.
- Kim, H.J., Chen, F., Wang, X., Rajapakse, N.C. (2005), Effect of chitosan on the biological properties of sweet basil (*Ocimum basilicum* L.), *J. Agric. Food Chem.*, 53, 3696-3701.
- Klimankova, E., Holadova, K., Hajslova, J., Cajka, T., Poustka, J., Koudela, M. (2008), Aroma profile of five basil (*Ocimum basilicum* L.) cultivars grown under conventional and organic conditions, *Food Chem.*, 107, 464-472.
- Koba, K., Poutouli, P.W., Raynaud, C., Chaumont, J., Sanda, K. (2009), Chemical composition and antimicrobial properties of different basil essential oils chemotypes from Togo. *Bangladesh. J. Pharmacol.*, 4, 1-8.
- Koche, D., Imran, S., Shirsat, R., Bhadange, D., (2011), Comparative phytochemical and nutritional studies of leaves and stem of three Lamiaceae members, *Rese. J. Pharm. Biol. Chem. Sci.*, 2, 1-4.
- Koroch, A.R., Juliani, H.R., Sims, C., Simon, J.E. (2010), Antioxidant activity, total phenolics and rosmarinic acid content in different basil (*Ocimum* spp.). *Isr. J. Plant Sci.*, 58, 191-195.
- Koyama, S., Purk, A., Kaur, M., Soini, H.A., Novotny, M.V., Davis, K., Kao, C.C., Matsunami, H., Mescher, A. (2019), Beta-caryophyllene enhances wound healing through multiple routes. *Plos One*, 14, e0216104.
- Krishan, G., Narang, A. (2014). Use of essential oils in poultry nutrition: A new approach. *J. Adv. Vet. Anim. Res.*, 1(4), 156-162.
- Krzysztof, C. (2006), Skin penetration of terpenes from essential oils and topical vehicles, *Planta Med.*, 72(4), 311-316.
- Kubiça, T.F., Alves, S.H., Weiblen, R., Lovato, L.T. (2014), *In vitro* inhibition of the bovine viral diarrhoea virus by the essential oil of *Ocimum basilicum* (basil) and monoterpenes, *Braz. J. Microbiol.*, 45(1), 209-214.
- Kumar, A., Agarwal, K., Maurya, A.K., Shanker, K., Bushra, U., Tandon, S. (2015). Pharmacological and phytochemical evaluation of *Ocimum sanctum* root extracts for it anti inflammatory, analgesic and antipyretic activities. *Phco.g Mag*, 11, 217-224.
- Kumar, A., Dubey, N.K, Srivastava, S., (2013), Antifungal evaluation of *Ocimum sanctum* essential oil against fungal deterioration of raw materials of *Rauvolfia serpentina* during storage, *Ind. Crops Prod.*, 45, 30-35.
- Laskar, S., Majumdar, S.G. (1988), Variation of major constituents of essential oil of the leaves of *Ocimum Sanctum* Linn, *J. Indian Chem. Soc.*, 65, 301-302.
- Lee, K.G., Shibamoto, T. (2001), Antioxidant property of aroma extract isolated from clove buds [*Syzygium aromaticum* (L.) Merr. Et Perry]. *Food Chem.*, 74(4), 443-448.
- Lee, S.J., Umamo, K., Shibamoto, T., Lee, K.G. (2005), Identification of volatile components in basil (*Ocimum basilicum* L.) and thyme leaves (*Thymus vulgaris* L.) and their antioxidant properties. *Food Chem.* 91, 131-137.
- Lee, J., Scagel, C.F. (2009). Chicoric acid found in basil (*Ocimum basilicum* L.) leaves. *Food Chem.*, 115, 650-656.
- Lee, J., Scagel, C.F. (2010). Chicoric acid levels in commercial basil (*Ocimum basilicum*) and *Echinacea purpurea* products. *J. Funct. Food.*, S2, 77 -84
- Letizia, C.S., Cocchiara, J., Lalko, J., Api, A.M. (2003), Fragrance material review on linalool. *Food Chem. Toxicol.*, 41, 943-964.
- Li, Y., Lai, Y., Wang, Y., Liu, Ni, Zhang, F., Xu, P. (2016). 1, 8-Cineole protect against influenza-virus-induced pneumonia in mice. *Inflammation*, 39, 1582-1593.
- Lokina, S., Stephen, A., Kaviyaran, V., Arulvasu, C., Narayanan, V. (2014), Cytotoxicity and antimicrobial activities of green synthesized silver nanoparticles, *Eur. J. Med. Chem.*, 76, 256-263.

- Lopez, M.D., Jordan, M.J., Pascual-Villalobos, M.J. (2008), Toxic compounds in essential oils of coriander, caraway and basil active against stored rice pests. *J. Stored Prod. Res.*, 44, 273-278.
- Losasso, C., Belluco, S., Cibir, V., Zavagnin, P., Mičetić, I., Gallochio, F., Zanella, M., Bregoli, L., Biancotto, G., Ricci, I. (2014), Antibacterial activity of silver nanoparticles: sensitivity of different *Salmonella* serovars. *Front. Microbiol.*, 5, 1-9.
- Loudon, I. (2006), A brief history of homeopathy, *J. R. Soc. Med.*, 99(12), 607-610.
- Machado, M.I.L., Silva, M.G.V., Matos, F.J.A., Craveiro, A.A., Alencar, J.W. (1999). Volatile constituents from leaves and inflorescence oil of *Ocimum tenuiflorum* L. f. (syn. *O. sanctum* L.) grown in Northeastern Brazil. *J. Essent. Oil Res.*, 11, 324-326.
- Malik, K., Arora, G., Singh, I. (2012). *Ocimum sanctum* seeds, a natural super disintegrant: Formulation and evaluation of fast melt tablets of nimesulide. *Polim Med.*, 42, 49-59. [This study shows that *Ocimum sanctum* seeds can be used as a natural superdisintegrant in the formulation of fast melt tablets].
- Mallikarjun, S., Rao, A., Rajesh, G., Shenoy, R., Pai, M. (2016), Antimicrobial efficacy of Tulsi leaf (*Ocimum sanctum*) extract on periodontal pathogens: An *in vitro* study. *J. Indian Soc. Periodontol.*, 20(2), 145-150.
- Manaharan, T., Thirugnanasampandan, R., Jayakumar, R., Kanthimathi, M.S., Ramya, G., Ramnath, M.G. (2016), Purified essential oil from *Ocimum sanctum* Linn. triggers the apoptotic mechanism in human breast cancer cells. *Phcog Mag.*, 12(46), 327-331.
- Manaharan, T., Thirugnanasampandan, R., Jayakumar, R., Ramya, G., Ramnath, G., Kanthimath, M.S. (2014), Antimetastatic and anti-inflammatory potentials of essential oil from edible *Ocimum sanctum* Leaves. *Sci. World J.*, ID 239508.
- Manikandan, P., Murugan, R.S., Abbas, H., Abraham, S.K., Nagini, S. (2007). *Ocimum sanctum* Linn. (Holy Basil) ethanolic leaf extract protects against 7, 12 dimethylbenz(a) anthracene induced genotoxicity, oxidative stress, and imbalance in xenobiotic metabolizing enzymes. *J. Med. Food*, 10, 495-502.
- Manzoor, F., Samreen, K.B., Parveen, Z. (2013). Larvicidal activity of essential oils against *Aedes aegypti* and *Culex quinquefasciatus* larvae (diptera: culicidae). *J. Anim. Plant Sci.*, 23, 420-424.
- Marotti, M., Piccaglia, R., Giovanelli, E. (1996), Differences in essential oil composition of basil (*Ocimum basilicum* L.) Italian cultivars related to morphological characteristics. *J. Agric. Food Chem.*, 14, 3926-3929.
- Martini, N., Eloff, J.N. (1998), The preliminary isolation of several antibacterial compounds from *Combretum erythrophyllum* (Combretaceae). *J. Ethnopharmacol.*, 62, 255-263.
- Medzhitov, R. (2010). Inflammation 2010: new adventures of an old flame. *Cell*, 140, 771-776.
- Mishra, M. (2008), Tulsi to Save Taj Mahal from Pollution Effects, *The Times of India*, Bennett Coleman and Co. Ltd.
- Mittal, R., Kumar, R., Chahal, H.S. (2018), Antimicrobial activity of *Ocimum sanctum* leaves extracts and oil. *J. Drug Deliv. Ther.*, 8(6), 201-204.
- Mondal, S., Mahapatra, S.C., Mirdha, B.R., Naik, S.N. (2007), Antimicrobial activity of essential oils obtained from fresh and dried leaves of *Ocimum sanctum* L. against enteric bacteria and yeast. *Acta Hortic.*, 756, 267-270.
- Mondawi, B.M., Duprey, R.J., Magboul, A.Z., Satti, A.M. (1984), Constituents of essential oil of *Ocimum basilicum* var. *thyrsiflorum*. *Fitoterapia*. 55, 60-61.
- Mosaddegh, M., Naghibi, F., Moazzeni, H., Pirani, A., Esmaeili, S. (2012). Ethnobotanical survey of herbal remedies traditionally used in kohgiluyehva Boyer Ahmad province of Iran. *J. Ethnopharmacol.*, 141, 80-95.
- Mousavi, L., Salleh, R.M., Murugaiyah, V. (2018). Phytochemical and bioactive compounds identification of *Ocimum tenuiflorum* leaves of methanol extract and its fraction with an anti-diabetic potential. *Int. J. Food Prop.*, 21(1), 2390-2399.

- Muenschner, C.W., Arthur, M. (1978), *Garden Spice and Wild Pot Herbs*, Ithaca, NY: Cornell University Press.
- Mukherji, S.P. (1995), Ocimum - a cheap source of Eugenol, *Science Reporter* 1987, pp. 599. 31. Eugenol-rich Ocimum variety released. In: *CSIR NEWS* (Published by PID CSIR, New Delhi); 45, 256.
- Muller, J., Greiner, J.F., Zeuner, M. (2016), 1,8-Cineole potentiates IFR3-mediated antiviral response in human stem cells and in ex vivo model of rhinosinusitis. *Clin. Sci. (Lond)*, 130(15), 1339-1359.
- Nahak, G., Mishra, R.C., Sahu, R.K. (2011), Phytochemical investigation and in vitro antioxidant evaluation of some *Ocimum* species. *J. Pharm.Res.*, 4(7), 2340-2343.
- Naidu, J.R., Ismail, R.B., Sasidharan S. (2016), Chemical profiling and antioxidant activity of Thai basil (*Ocimum basilicum*). *J. Essent. Oil Bear. Pl.*, 19(3), 750-755.
- Nakamura, T.U., Mendonça, R.R., Morgado, J.A., Maza, P.K., Dias, B.P., García, D.A., Alviano, D.S., do, M., Rosa, S., Lopes, A.H., Alviano, C.S., Nakamura, C.V. (2006), Antileishmanial activity of eugenol-rich essential oil from *Ocimum gratissimum*. *Int. J. Parasitol.*, 55, 99-105.
- Narusuye, K., Kawai, F., Matsuzaki, K., Miyachi, E. (2005), Linalool suppresses voltage-gated currents in sensory neurons and cerebellar Purkinje cells. *J. Neural Transm.*, 112, 193-203.
- Nascimento, L.D., de Moraes, A.A.B., da Costa, K.S., Galúcio, J.M.P., Taube, P.S., Costa, C.M.L., Cruz, J.N., Andrade, E.H.A., de Faria, L.J.G. (2020). Bioactive natural compounds and antioxidant activity of essential oils from spice plants: new findings and potential applications. *Biomolecules*, 10(7), 988.
- Nathan, C., Ding, A. (2010), Nonresolving inflammation. *Cell*, 140, 871-882.
- Negahban, M., Saeedfar, S., Zakerin, A., Aboutalebi, A. (2015), The effect of different harvest stages on the quality and quantity of the essential oil of tulsi (*Ocimum sanctum* L.). *Russ. J. Biol Res.*, 3(1), 39-42.
- Nishijima, H., Uchida, R., Kimiko, K., Kawakami, N., Ohkuba, T., Kitamura, K. (1999), Mechanisms mediating the vasorelaxing action of eugenol, pungent oil, on rabbit arterial tissues, *Jap. J. Pharmacol.*, 79(3), 327-334.
- Nour, A.H., Nour, A.H., Yusoff, M.M., Jessinta, S. (2012). Bioactive compounds from basil (*Ocimum basilicum*) essential oils with larvicidal activity against *Aedes aegypti* larvae. 3rd International Conference on Biology, Environment and Chemistry IPCBEE;
- Núñez, I.C., Mendonça, P.M., Arranz, J.C.E., Cortinhas, L.B., Dutok-Sánchez, C.M., Queiroz, M.M.C. (2018). Insecticidal effects of *Ocimum sanctum* var. *cubensis* essential oil on the diseases vector *Chrysomya putoria*. *J. Pharm. Pharmacogn. Res.*, 6 (3), 148-157.
- Okoye, F.B.C., Obonga, W.O., Onyegbule, F.A., Ndu, O.O., Ihekwereme, C.P. (2014), Chemical composition and anti-inflammatory activity of essential oils from the leaves of *Ocimum basilicum* L. and *Ocimum gratissimum* L. (Lamiaceae), *Int. J. Pharm. Sci. Res.*, 5(6), 2174-2180.
- Oliveira, J.S., Porto, L.A., Estevam, C.S., Siqueira, R.S., Alves, P.B., Niculau, E.S. (2009). Phytochemical screening and anticonvulsant property of *Ocimum basilicum* leaf essential oil. *B. Latinoam. Caribe Pl.*, 8, 195-202.
- Ozcan, M., Chalchat, J. (2002), Essential oil composition of *Ocimum basilicum* L. and *Ocimum minimum* L. in Turkey, *Czech J. Food Sci.*, 20, 223-228.
- Pachkore, G.L., Dhale, D.A. (2012). Phytochemicals, vitamins and minerals content of three *Ocimum* species. *International Journal of Science Innovations and Discoveries*, 2, 201-207.
- Padalia, R.C., Verma, R.S. (2011), Comparative volatile oil composition of four *Ocimum* species from northern India. *Nat. Prod. Res.*, 25(6), 569-575.
- Pandey, A.K., Tripathi, N.N. (2011), Post-harvest fungal and insect deterioration of pigeon pea seeds and their management by plant volatiles, *J. Indian Bot.Soc.*, 90, 326-331.
- Panneerselvam, C., Murugan, K., Amerasan, D. (2015), Biosynthesis of silver nanoparticles using plant extract and its anti-plasmodial property, *Adv. Mat. Res.*, 1086, 11-30.

- Parag, S., Vijayashree, N., Rami, B., Patil, B. (2010), Antibacterial activity of *Ocimum sanctum* Linn. and its application in water purification, *Res. J. Chem. Environ.*, 14, 46-50.
- Pathak, I., Niraula, M. (2019), Assessment of total phenolic, flavonoid content and antioxidant activity of *Ocimum sanctum* Linn., *J. Nep. Chem. Soc.*, 40, 30-35.
- Patil, R., Patil, R., Ahirwar, B., Ahiwar, D. (2011), Isolation and characterization of anti-diabetic component (bioactivity-guided fractionation) from *Ocimum sanctum* L. (Lamiaceae) aerial part, *Asian Pac. J. Trop. Med.*, 4(4), 278-282.
- Pattanayak, P., Behera, P., Das, D., Panda, S.K. (2010). A reservoir plant for therapeutic applications: an overview. *Pharmacogn. Rev.*, 4, 95-105.
- Pavela, R., Kaffkova, K., Kumsta, M. (2014), Chemical composition and larvicidal activity of essential oils from different *Mentha* L. and *Pulegium* species against *Culex quinquefasciatus* say (Diptera: Culicidae). *Plant Prot. Sci.*, 50(1), 36-42.
- Pawar, V.C., Thaker, V.S. (2006). In vitro efficacy of 75 essential oils against *Aspergillus niger*. *Mycoses*, 49, 316-323.
- Peana, A.T., De Montis, V.I.G., Nieddu, L., Spano, M.T., D'Aquila, P.S., Pippia, P. (2004a). Profile of spinal and supra-spinal antinociception or (-)-linalool. *Eur. J. Pharmacol.*, 485, 165-174.
- Peana, A.T., D'Aquila, P.S., Chessa, M.L., Moretti, M.D.L., Serra, G., Pippia, P. (2003), (-)-Linalool produces antinociception in two experimental models of pain. *Eur. J. Pharmacol.*, 460, 37-41.
- Peana, A.T., D'Aquila, P.S., Panin, F., Serra, G., Pippia, P., Moretti, M.D.L. (2002), Anti-inflammatory activity of linalool and linalyl acetate constituents of essential oils, *Phytomedicine*, 9(8), 721-726.
- Pemminati, S., Gopalakrishna, H.N., Venkatesh, V., Rai, A., Shetty, S., Vinod, A. (2011), Anxiolytic effect of acute administration of ursolic acid in rats, *Res. J. Pharm. Biol. Chem. Sci.*, 2, 431-437.
- Petrovic, J., Stojković, D., Sokovic, M. (2019), Terpene core in selected aromatic and edible plants: Natural health improving agents. *Adv. Food Nutr. Res.*, 90, 423-451.
- Pierik, R., Ballare, C.L., Dicke, M. (2014), Ecology of plant volatiles: taking a plant community perspective. *Plant Cell Environ.*, 37, 1845-1853.
- Pingale, S.S., Firke N.P.B, Markandeya, A.G. (2012). Therapeutic activities of *Ocimum tenuiflorum* accounted in last decade: A review. *J. Pharm. Res.*, 5(4), 2215-2220.
- Pino, J.A., Rosado, A., Rodriguez, M., Garcia, D. (1998), Composition of the essential oil of *Ocimum tenuiflorum* L. grown in Cuba, *J. Essent. Oil Res.*, 10, 437-438.
- Plócica, J., Tal-Figiel, B. (2011). Selected plant and essential oils used in cosmetic industry. *Polish Journal of Cosmetology*, 14(4), 266-272.
- Politeo, O., Jukic, M., Milos, M. (2007), Chemical composition and antioxidant capacity of free volatile aglycones from basil (*Ocimum basilicum* L.) compared with its essential oil. *Food Chem.* 101, 379-385.
- Prabhu, D., Arulvasu, C., Babu, G., Manikandan, R., Srinivasan, P. (2013). Biologically synthesized green silver nanoparticles from leaf extract of *Vitex negundo* L. induce growth-inhibitory effect on human colon cancer cell line HCT15. *Process Biochem.*, 48, 317-324.
- Prakash, J., Gupta, S.K. (2000), Chemopreventive activity of *Ocimum sanctum* seed oil, *J Ethnopharmacol.* 72, 29-34.
- Prakash, J., Gupta, S.K. (2005). Therapeutic uses of *Ocimum sanctum* Linn (tulsi) with a note on eugenol and its pharmacological actions: a short review. *Indian J. Physiol. Pharmacol.*, 49, 125-131.
- Prakash, P., Neelu, G. (2005), Therapeutic uses of *Ocimum sanctum* linn (tulsi) with a note on eugenol and its pharmacological actions, *Indian J. Physiol. Pharmacol.*, 49(2), 125-131 [This review has discussed the role of *Ocimum sanctum* and its main constituent eugenol in the management of various ailments].

Prashar, R., Kumar, A., Hewer, A., Cole, K.J., Davis, W. (1998). Phillips DH. Inhibition by and extract of *Ocimum sanctum* of DNA binding activity of 7,12 dimethylbenz[a] anthracene in rat hepatocytes in vitro. *Cancer Lett.*, 128, 155-160.

chemical Pushpangadan, P., George, V., (2012), *Basil. Handbook of Herbs and Species*, Woodhead publishing limited.

Qamar, K.S., Dar, A.S., Bina, S., Kabir, N., Aslam, H. (2010), Anticancer activity of *Ocimum basilicum* and the effect of ursolic acid on the cytoskeleton of MCF-7 human breast cancer cells, *Lett. Drug Des. Discov.*, 7, 726-736.

Qi, Li, Li, Hai-Jiao, Xu, Tong, Du, Huan, Gang, Chen-Lei Huan, Fan, Gang, Zhang, Yi. (2018), Natural medicines used in the traditional tibetan medical system for the treatment of liver diseases, *Front. Pharmacol.*, 9(29). [In this study, traditional Tibetan medicine and its utilization to control the liver diseases have been discussed].

Quintans, J.S.S., Shanmugaam, S., Heimfarth, L. (2018). Monoterpenes modulating cytokines: a review. *Food Chem. Toxicol.*, 123, 233-257. [This study explores the antiinflammatory potential of monoterpenes].

Raghavendra, M., Maiti, R., Kumar, S., Acharya, S.B. (2009), Role of *Ocimum sanctum* in the experimental model of Alzheimer's disease in rats, *Int. J. Green Pharm.*, 3, 6-15.

Raguso, R.C., Pichersky, E. (1999). New Perspectives in Pollination Biology: Floral Fragrances. A day in the life of a linalool molecule: Chemical communication in a plant-pollinator system. Part 1: Linalool biosynthesis in flowering plants. *Plant Species Biol.*, 14, 95-120.

Raina, P., Deepak, M., Chandrasekaran, C.V., Agarwal, A., Wagh, N., Kaul-Ghanekar, R. (2016), Comparative analysis of anti-inflammatory activity of aqueous and methanolic extracts of *Ocimum basilicum* in RAW264.7, SW1353 and human primary chondrocytes. *J. Herb. Med.*, 6, 28-36.

Rajamma, A.J., Dubey, S., Sateesha, S.B., Tiwari, S.N., Ghosh S.K. (2011), Comparative larvicidal activity of different species of *Ocimum* against *Culex Quinquefasciatus*, *J. Nat. Prod.*, 25(20), 1916-1922.

Rakha, P., Sharma, S. and Parle, M. (2010), Anti-inflammatory potential of the seeds of *Ocimum basilicum* Linn. in rats, *Asian J. Biol. Sci.*, 5, 16-18.

Rameshrad, M., Salehian, R., Fathiazad, F., Hamedeyazdan, S., Garjani, M., Maleki-Dizaji, N. (2014). The effects of *Ocimum basilicum* ethanol extract on carrageenan induced paw inflammation in rats. *Pharm. Sci.*, 20, 149-156.

Rana, L., Tewari, G., Pande, C., (2020). Phytochemical and pharmacological overview on *Ocimum sanctum* linn.: effect of growth stages, In: Natural Products and Their Utilization Pattern (Eds. G. Tewari, A. Tewari, A. Tewari, L. M. Tewari). Nova Science Publishers, Inc.

Rates, S.M.K. (2001), Plants as source of drugs, *Toxicon*, 39, 603-613.

Rattan, R.S. (2010). Mechanism of action of insecticidal secondary metabolites of plant origin, *J. Crop Prot.*, 29, 913-920.

Rattanachaikunsopon, P., Phumkhachorn, P. (2010), Antimicrobial activity of basil (*Ocimum basilicum*) oil against salmonella enteritidis in vitro and in food. *Biosci. Biotechnol. Biochem.*, 74(6), 1200-1204.

Rezzoug, M., Bakchiche, B., Gherib, A., Roberta, A., Guido, F., Kilinçarslan, O., Mammadov, R., Bardaweel, S.K. (2019), Chemical composition and bioactivity of essential oils and ethanolic extracts of *Ocimum basilicum* L. and *Thymus algeriensis* Boiss. and Reut. from the Algerian Saharan Atlas. *BMC Complement. Altern. Med.*, 19, 146-156.

Rizwana, K. (2018), Antifungal activity test by aqueous solution of *Ocimum sanctum* [Tulsi]. *Int. J. Adv. Sci. Eng. Technol.*, 6 (4).

Rodrigues, L.B., Martins, A.O.B.P.B., Cesário, F.R., Castro, F.E.F., de Albuquerque, T.R., Fernandes, M.N.M., Silva, F.B.A., Júnior, L.J.Q., da Costa, J.G., Coutinho, H.D.M., Barbosa, R., de Menezes, I.R.A. (2016). Anti-inflammatory and antiedematogenic activity of the *Ocimum basilicum* essential oil and its main compound estragole: *in vivo* mouse models. *Chem. Biol. Interact.*, 257, 4-25.

- Rodríguez-González, Á., Álvarez-García, S., González-López, O., Silva, F.D., Casquero, P.A. (2019), Insecticidal properties of *Ocimum basilicum* and *Cymbopogon winterianus* against *Acanthoscelides obtectus*, insect pest of the common bean (*Phaseolus vulgaris*, L.). *Insects*, 10(5), 151.
- Sá, R.D.C.D.S.E., Lima, T.C., Da Nóbrega, F.R., De Brito, A.E.M., De Sousa, D.P. (2017). Analgesic-like activity of essential oil constituents: an update. *Int. J. Mol. Sci.*, 18, 2392.
- Saaban, K.F., Ang, C.H., Khor, S.M., Chuah, C.H. (2019), Chemical Constituents and antioxidant capacity of *Ocimum basilicum* and *Ocimum sanctum* Iran. *J. Chem. Chem. Eng.* 38(2), 139-152.
- Saha, S., Dhar, T.N., Sengupta, C., Ghosh, P. (2013), Biological activities of essential oils and methanol extracts of five *Ocimum* species against pathogenic bacteria, *Czech J. Food Sci.*, 31, 194–202.
- Saha, S., Mukhopadhyay, M.K., Ghosh, P.D., Nath, D. (2012), Effect of methanolic leaf extract of *Ocimum basilicum* L. on benzene-induced hematotoxicity in mice, *Evid. Based Complement. Alternat. Med.*, (3), 176385.
- Sailaja, I., Shaker A.I. (2010), Antioxidant activity in *Ocimum sanctum* Linn., *Ocimum basilicum*. *Asian J. Biol. Sci.*, 5(2), 195-199.
- Sani, Y.A., Cyril, O., Yahaya, Y., Hassan, U., Ibrahim, L.S. (2018), Phytochemical screening, in vitro antibacterial and partial TLC purification of different solvents extracts of *Ocimum basilicum* L. *J. Pharm. Biol. Sci.* 5(02), 132-138.
- Santoro, G.F., Cardoso, M.G., Guimaraes, L.G., Mendonca, L.Z., Soares, M.J. (2007). Trypanosoma cruzi: activity of essential oils from *Achillea millefolium* L., *Syzygium aromaticum* L. and *Ocimum basilicum* L. on epimastigotes and trypomastigotes. *Exp. Parasitol.*, 116, 283-290.
- Santos, F.A., Rao, V.S. (2000), Anti-inflammatory and antinociceptive effects of 1, 8-cineole, a terpenoid oxide present in many plant essential oils, *Photother. Res.*, 4, 240-244.
- Sarahroodi, S., Esmaeili, S., Mikaili, P., Hemmati, Z., Saberi, Y. (2012), The effect of green *Ocimum basilicum* hydroalcoholic extract on retention and retrieval of memory in mice. *Anc. Sci. Life*, 31(4), 185-189.
- Sarkar, A., Pandey, D.N., Pant, M.C. (1990), A report on the effect of *Ocimum sanctum* (Tulsi) leaves and seeds on blood and urinary uric acid, urea and urine volume in normal albino rabbits, *Indian J. Physiol. Pharmacol.*, 34, 61-62.
- Sastry, K.P., Kumar, R.R., Kumar, A.N., Sneha, G., Elizabeth, M. (2012), Morpho chemical description and antimicrobial activity of different *Ocimum* species, *J. Plant. Dev.*, 19, 53-64.
- Schott, M., Klein, B., Vilcinskis, A. (2015), Detection of illicit drugs by trained honeybees (*Apis mellifera*). *Plos One*, 10(6), e0128528.
- Scriven, R., Meloan, C.E. (1984), Determining the active component in 1.3.3-Trimethyl-2-oxabicyclo [2.2.2] octane (cineole) that repel the American cockroach, *Periplaneta Americana*. *Ohio J. Sci.*, 84(3), 85-88.
- Selvakkumar, C., Gayathri, B., Vinaykumar, K.S., Lakshmi, B.S., Balakrishnan, A. (2007), Potential antiinflammatory properties of crude alcoholic extract of *Ocimum basilicum* L. in human peripheral blood mononuclear cells. *J. Health Sci.*, 53, 500-505.
- Sen, P. (1993). Therapeutic potentials of Tulsi: from experience to facts, *Drugs News & Views*, 1(2), 15-21.
- Sharafati-Chaleshtori R, Rokni N, Rafieian-Kopaei M, Drees F, Salehi E. (2015), Antioxidant and antibacterial activity of basil (*Ocimum basilicum* L.) essential oil in beef burger, *J. Agric. Sci. Technol.* 17, 817-826.
- Sharma, S., Singh, R., Thakre, B. (2019), Antifungal activity of leaf extracts of *Ocimum sanctum* against fungal pathogens. *Int. J. Curr. Microbiol. App. Sci.*, 8(4), 1210-1214.
- Sharopov, F.S., Satyal, P., Ali, N.A.A., Pokharel, S., Zhang, H., Wink, M., Kukaniev, M.A., Setzer, W.N. (2016). The essential oil compositions of *Ocimum basilicum* from three different regions: Nepal, Tajikistan, and Yemen. *Chem. Biodivers.*, 13(2), 241-248.

- Sheoran, N., Kumar, R., Kumar, A., Batra, K., Sihag, S., Maan, S., Maan, N.S. (2017), Nutrigenomic evaluation of garlic (*Allium sativum*) and holy basil (*Ocimum sanctum*) leaf powder supplementation on growth performance and immune characteristics in broilers. *Vet. World*, 10(1), 121-129.
- Shirazi, M.T., Gholami, H., Kavooosi, G., Rowshan, V., Tafsiy, A. (2014), Chemical composition, antioxidant, antimicrobial and cytotoxic activities of *Tagetes minuta* and *Ocimum basilicum* essential oils, *Food Sci. Nutr.*, 2(2), 146-155.
- Shivananjappa, M. Joshi, M. (2012), Aqueous extract of tulsi (*Ocimum sanctum*) enhances endogenous antioxidant defenses of human hepatoma cell line (HepG2), *J. Herbs Spices Med. Plants*, 18(4), 331-348.
- Sienkiewicz, M., Lysakowska, M., Pastuszka, M., Bienias, W., Kowalczyk, E. (2013), The potential use of basil and rosemary essential oils as effective antibacterial agents, *Molecules*, 18, 9334-9351.
- Silva, A.V., Sousa, J.P., Pessôa, H.L.F., Freitas, A.F.R., Coutinho, H.D.M., Alves, L.B.N., Lima, E.O. (2016), *Ocimum basilicum*: Antibacterial activity and association study with antibiotics against bacteria of clinical importance. *Pharm Biol.*, 54(5), 863-867.
- Silva, L., Emanuelli, T., Souza, D., Elisabetsky, E. (2001), Effects of linalool on glutamate release and uptake in mouse cortical synaptosomes. *Neurochem. Res.*, 26, 191-194.
- Sims, C.A., Juliani, H.R., Mentreddy, S.R. and Simon, J.E. (2014), Agronom, essential oils in holy basil (*Ocimum tenuiflorum* L.) as influenced by planting dates and harvest times in North Alabama. *Journal of Medicinally Active Plants.*, 2(3-4), 33-41.
- Singh, D., Chaudhuri, P.K. (2018), A review on phytochemical and pharmacological properties of Holy basil (*Ocimum sanctum* L.), *Ind. Crops Prod.*, 118, 367-382.
- Singh, P., Mittal, V.K., Gupta, S.C. (2003), Trace elements in typical herbs as an indicator of environmental pollution, *Indian J. Environ. Prot.*, 23, 1114-1119.
- Singh, S., Mahour, K., Prakash S. (2009), Evaluation of mosquito repellent efficacy of *Ocimum sanctum* plant extract. *J. Herb Med. Toxicol.*, 3, 87-90
- Singh, S., Majumdar, D.K., Rehan, H.M.S. (1996), Evaluation of anti-inflammatory potential of fixed oil of *Ocimum sanctum* (Holybasil) and its possible mechanism of action. *J. Ethnopharmacol.*, 54(1), 19-26.
- Singh, S., Malhotra, M., Majumdar, D.K. (2005), Antibacterial activity of *Ocimum sanctum* L. fixed oil, *Indian J. Exp. Biol.*, 43, 835-837.
- Singh, S., Taneja, M., Majumdar, D.K. (2007), Biological activities of *Ocimum sanctum* L. fixed oil--an overview, *Indian J. Exp. Biol.*, 45, 403-412.
- Smitha, G.R., Varghese, T.S., Manivel, P. (2016). *Cultivation of Ocimum*. Extension Bulletin, Anand Press, Anand, Gujarat,
- Soni, A., Sosa, S., (2013), Phytochemical analysis and free radical scavenging potential of herbal and medicinal plant extracts, *Int. J. Pharmacogn. Phytochem.*, 2 (4), 22-29.
- Sousa, O.V., Silverio, M.S., Del-Vechio-Vieira, G., Matheus, F.C., Yamamoto, C.H., Alves, M.S. (2008), Antinociceptive and anti-inflammatory effects of the essential oil from *Eremanthus erythropappus* leaves, *J. Pharm. Pharmacol.*, 60, 771-777.
- Srivastava, P., Singh, M., Devi, G., Chaturvedi, R. (2020), Herbal medicine and biotechnology for the benefit of human health animal biotechnology: Applications and concerns. In: *Animal Biotechnology*. 2nd Edition. (Eds. A.S.Verma, A. Singh). Academic Press. pp. 613-629.
- Suanarunsawat, T., Anantasomboon, G., Piewban, C. (2016), Anti-diabetic and anti-oxidative activity of fixed oil extracted from *Ocimum sanctum* L. leaves in diabetic rats. *Exp. Ther. Med.*, 11(3), 832-840.
- Subramanian, M., Chintalwar, G.J., Chattopadhyay S. (2005), Antioxidant and radioprotective properties of an *Ocimum sanctum* polysaccharide, *Redox Rep.*, 10, 257-264.
- Sundaram, R.S., Gowtham, L., Ramanathan, M., Manikandan, P., Venugopal, V., Kamalakannan, D., Nayak, B.S. (2011), Quantification of bioactive principles in Indian traditional herb *Ocimum sanctum* Linn., (Holy Basil) leaves by high performance liquid chromatography. *Asian J. Pharm. Biol. Sci.*, 1 (3), 35-41.

- Sundaramurthi, P., Dhandapani, S., Ponnusamy, S., Subbaiyan, M. (2012), Effect of tulsi (*Ocimum sanctum*) as a disinfectant for water treatment, *J. Biosci. Bioeng.*, 1(1), 1-7.
- Sundarraju, D., Anbu, J., Reeta, R., Senthilkumar, K.L., Anjana, A. (2014): Pharmacognostical and phytochemical investigation of ethanolic extract of *Ocimum basilicum* Linn. *Int. J. Pharm. Chem. Biol. Sci.*, 4(1), 194-200.
- Suthar, S. K., Malik, A. K. (2015), GC/MS analysis of volatile compounds of the essential oil of leaves of *Ocimum sanctum* growing in Hisar, India. *Asian J. Chem.* 27(8), 3135-3136.
- Tajkarimi, M.M., Ibrahim, S.A., Cliver, D.O. (2010), Antimicrobial herb and spice compounds in food, *Food Control*, 21, 1199-1218. [This study has convincingly demonstrated about the antimicrobial activity of spice and aromatic herbs which are used in food materials].
- Tangpao, T., Chung, H.H., Sommano, S.R. (2018), Aromatic profiles of essential oils from five commonly used Thai basil. *Foods*, 7(11), 175.
- Tapondjou, L.A., Adler, C., Bouda, H., Fontem, D.A. (2002). Efficacy of powder and essential oil from *Chenopodium ambrosioides* leaves as post-harvest grain protectants against six-stored product beetles. *J. Stored Prod. Res.*, 38, 395-402.
- Telci, I., Bayram, E., Yilmaz, G., Avci, B. (2006), Variability in essential oil composition of Turkish basil (*Ocimum basilicum* L.). *Biochem. Syst. Ecol.*, 34, 489-497.
- Thiele, J.J., Dreher, F., Packer, L. (2000), Antioxidant defense systems in skin, in: P Elsner, HI Maibach, eds., *Cosmeceuticals, Drugs versus Cosmetics*, New York: Marcel Dekker 145-187 pp. Tomar, U.S., Daniel, V., Shrivastava, K., Panwar, M.S., Pant P. (2010). Comparative evaluation and antimicrobial activity of *Ocimum basilicum* Linn. (Labiatae) *J. Global Pharmacol. Technol.* 2, 49-53.
- Trettel, J.R., Gazim, Z.C., Goncalves, J.E., Stracieri, J., Magalhaes, H.M. (2017), Volatile essential oil chemical composition of basil (*Ocimum basilicum* L. 'Green') cultivated in a greenhouse and micropropagated on a culture medium containing copper sulfate. *In Vitro Cell. Dev. Biol. Plant.*, 53, 631-640.
- Unnithan, C.R., Dagnaw, W., Undrala, S., Ravi, S. (2013), Chemical composition and antibacterial activity of essential oil of *Ocimum basilicum* of Northern Ethiopia, *Int. Res. J. Biol. Sci.*, 2, 1-4.
- Valente, J., Zuzarte, M., Goncalves, M.J., Lopes, M.C., Cavaleiro, C., Salgueiro, L., Cruz, M.T. (2013). Antifungal, antioxidant and anti-inflammatory activities of *Oenanthe crocata* L. essential oil, *Food Chem. Toxicol.*, 62, 349-354.
- Vani, S.R., Cheng, S.F., Chuah, S.H. (2009), Comparative study of volatile compounds from genus *Ocimum*. *Am. J. Appl. Sci.*, 6 (3), 523-528. [This study elaborates the compositional variation of genus *Ocimum*].
- Vidhani, S.I., Vyas, V.G., Parmar, H.J., Bhalani, V.M., Hasan, M.M., Gaber, A., Golakiya, B.A. (2016). Evaluation of some chemical composition, minerals fatty acid profiles, antioxidant and antimicrobial activities of Tulsi (*Ocimum sanctum*) from India. *Am. J. Food Sci. Technol.*, 4, 52-57.
- Vlase, L., Benedec, D., Hanganu, D., Damian, G., Csillag, I., Sevastre, B. (2014). Evaluation of antioxidant and antimicrobial activities and phenolic profile for *Hyssopus officinalis* and *Teucrium chamaedrys*. *Molecules*, 19(5), 5490-5507.
- Wani, N.S., Bhalerao, A.K., Ranaware, V.P., Zanje, R. (2013), Formulation and evaluation of herbal sanitizer, *Int. J. Pharmtech Res.*, 5, 40-43. [This study formulated herbal sanitizer from aromatic plants and its biological effects].
- Warsi, S., Sholichah, A.R. (2017). A. Phytochemical screening and antioxidant activity of ethanolic extract and ethyl acetate fraction from basil leaf (*Ocimum basilicum* L.) by DPPH radical scavenging method. *IOP Conf. Ser.: Mater. Sci. Eng.*, 259:012008.
- Worth, H., Schacher, C., Dethlefsen, U. (2009), Concomitant therapy with cineole (eucalyptole) reduces exacerbations in COPD: a placebo-controlled double-blind trial. *Respir. Res.*, 10, 69. [This study explains the importance of cineole as an active controller of airway inflammation]

- Xia, K.Z., Perveen, N., Khan, N.H. (2018), Phytochemical analysis, antibacterial and antioxidant activity determination of *Ocimum sanctum*, *Pharm. Pharmacol. Int. J.*, 6(6).
- Xiao, M., Liu, R., Long, C., Ruan, Y., Liu, C. (2020). Using β -ocimene to increase the artemisinin content in juvenile plants of *Artemisia annua* L. *Biotechnol. Lett.*, 42, 1161-1167 [In this paper, the authors summried the use of β -ocimene to enhance artemisinin accumulation in juvenile *A. annua* plants]
- Yadav, N.P., Khatri, R., Bawankule, D.U., Pal, A., Shanker, K., Srivastava, P. (2009). Topical anti-inflammatory effects of *Ocimum basilicum* leaf extract in the phorbol-12, 13-dibutyrate model of mouse ear inflammation. *Planta Med.*, 75-PA72.
- Yamani, H.A., Pang, E.C., Mantri, N., Deighton, M.A. (2016), Antimicrobial activity of tulsi (*Ocimum tenuiflorum*) essential oil and their major constituents against three species of bacteria. *Front. Microbiol.* 7.
- Youssef, D.A., El-Fayoumi, H.M., Mahmoud, M.F. (2019), Beta-caryophyllene protects against diet-induced dyslipidemia and vascular inflammation in rats: Involvement of CB2 and PPAR-gamma receptors. *Chem-Biol Interact.*, 297, 16-24.
- Zafar, I., Muhammad, K.S., Khurram, S., Shaista, N. (2018), Antioxidant activity of essential oil from *Lawsoniainermis* Linn from Pakistan, *Int. J. Biosci.*, 12(3), 110-115.
- Zheljaskov, V.D., Cantrell, C.L., Evans, W.B., Ebelhar, M.W., Coker, C. (2008b), Yield and composition of *Ocimum basilicum* L. and *Ocimum sanctum* L. grown at four locations, *Hort. Sci.* 43, 737-741.
- Zheljaskov, V.D., Cantrell, C.L., Tekwani, B., Khan, S.I. (2008a), Content, composition, and bioactivity of the essential oils of three basil genotypes as a function of harvesting, *J. Agric. Food Chem.*, 56, 380-385. [Proposed studied data suggested the effect of harvest period on essential oil content, profile and bioactivities]
- Złoteka, U., Mikulskaa, S., Nagajeka, M., Wieca, M. (2016), The effect of different solvents and number of extraction steps on the polyphenol content and antioxidant capacity of basil leaves (*Ocimum basilicum*L.) extracts. *Saudi J. Biol. Sci.*, 23(5), 628-633.
- Zollo, P.H., Biyiti, L., Tchoumboungang, F., Menut, C., Lamaty, C., Bouchet, P.H. (1998), Aromatic plants of tropical Africa. Part XXXII. Chemical composition and antifungal activity of thirteen essential oils from aromatic plants of Cameroon, *Flavour. Frag. J.*, 13, 107-114.

Biographical Sketches

Lata Rana received her master's degree in chemistry (2016) from Kumaun University, Nainital, Uttarakhand, India. At present, her doctoral programme is ongoing from Kumaun University. She has 7 research publications, 2 book chapters and participated more than 20 conferences/ workshops/ webinars.

Dr. Geeta Tewari, M. Sc., , Ph. D., is working as Associate Professor in the Department of Chemistry, D. S. B. Campus, Kumaun University, Nainital. She has authored two text books and two edited books. She has published more than 75 research papers 19 book chapters and more than 20 popular articles in different magazines.

Dr. Deepshikha Punetha received her Ph.D. degree from Kumaun University Nainital, Uttarakhand, India. She has published 10 research papers and 4 book chapters. Currently, she is working in Delma Laboratory for Soil & Construction Material Inspection L.L.C, Abu Dhabi, United Arab Emirates.

Dr. Mamta Bisht received her Ph. D. degree in Natural product research from D. S. B. Campus, Kumaun University, Nainital, India. She has published more than ten research papers and one book chapter.

Geetanjali Upadhyay did her master's degree in Botany (2018) from Kumaun University Nainital, Uttarakhand, India. Currently, she is persuing Ph. D. from Kumaun University, Nainital.

Dr. Chitra Pande, M. Sc., Ph. D., is working as Professor in the Department of Chemistry, D. S. B. Campus, Kumaun University, Nainital. She has published 6 bookd chapters and more than 60 research papers in various national and international journals.

Dr. Lalit M. Tewari, M. Sc., Ph. D., is working as Professor in the Department of Botany, D. S. B. Campus, Nainital and Director, Research & Extension, Kumaun University. He has authored ten Edited

MEDICINAL AND AROMATIC PLANTS OF THE WORLD - *Therapeutic Potential and Phytochemistry of Natural Herbals of Family Lamiaceae: Ocimum Sanctum L. and Ocimum Basilicum* - Lata Rana, Geeta Tewari, Deepshikha Punetha, Mamta Bisht, Geetanjali Upadhyay, Chitra Pande and Lalit M. Tewari

Magazines on Uttarakhand culture and fifteen books. He has published more than 145 research papers, 10 book chapters and more than 50 popular articles in different magazines.